

**Washington County** 2016 Shared Community Health Needs Assessment

# Acknowledgements

The following report is funded through the generous support and contributions of the Maine Shared Health Needs Assessment Planning Process Collaborative:



The report was prepared by the research teams at Market Decisions Research of Portland, Maine, Hart Consulting Inc. of Gardiner, Maine, and the Maine Center for Disease Control and Prevention. Substantial segments of the narrative sections were adapted from the 2012 Maine State Health Assessment and significant analysis and research was conducted by epidemiologists at the Maine CDC and the University of Southern Maine's Muskie School of Public Service.



See end of the report for a list of contributors and collaborating organizations.

Maine Shared Community Health Needs Assessment, 2015 © 2015 MaineGeneral Medical Center, subject to perpetual rights of use by Eastern Maine Healthcare Systems, MaineHealth, Central Maine Healthcare and Maine Center for Disease Control and Prevention, an office of the Department of Health and Human Services.

Note: Originally, this report was dated 2015 on the cover. However, it has been changed to 2016 to reflect the fiscal years of the organizations that have been involved.

# Table of Contents

Table of Tables	iii
How to Use This Report	v
Executive Summary	1
Background	
County Demographics	
Washington County Summary of Findings	9
Washington County Priority Health Issues and Factors	
County Health Rankings & Roadmaps	
Stakeholder Survey Findings	
Health Indicators Results from Secondary Data Sources	41

This page is blank intentionally.

# Table of Tables

Table 1. Key Socioeconomic Indicators for Washington County
Table 2. Key Health and Mortality Indicators for Washington County
Table 3. Key Access to Health/Health Care Quality Indicators for Washington County 11
Table 4. Key Asthma and COPD Indicators for Washington County
Table 5. Key Cancer Indicators for Washington County    12
Table 6. Key Cardiovascular Disease Indicators for Washington County
Table 7. Key Diabetes Indicators for Washington County    13
Table 8. Key Environmental Health Indicators for Washington County
Table 9. Key Immunization Indicators for Washington County
Table 10. Key Infectious Disease Indicators for Washington County
Table 11. Key Sexually Transmitted Disease Indicators for Washington County
Table 12. Key Intentional Injury Indicators for Washington County
Table 13. Key Unintentional Injury Indicators for Washington County         16
Table 14. Key Mental Health Indicators for Washington County    17
Table 15. Key Nutrition and Physical Activity Indicators for Washington County
Table 16. Key Weight Indicators for Washington County    19
Table 17. Key Pregnancy and Birth Outcomes for Washington County
Table 18. Key Substance Abuse Indicators for Washington County
Table 19. Key Tobacco Use Indicators for Washington County    22
Table 20. Percentage of Stakeholders who agreed that Significant Disparities Exist AmongSpecific Groups for a Specific Health Issue
Table 21. Percentage of Stakeholders who identified Certain Factors as Key Drivers thatlead to a Specific Health Condition
Table 22. Priority Health Issue Successes and Challenges for Washington County-Surveillance Data29
Table 23. Priority Health Issue Challenges and Resources for Washington County-Stakeholder Survey Responses32
Table 24. Priority Health Factor Strengths and Challenges for Washington County-Surveillance Data33
Table 25. Priority Health Factor Challenges and Resources for Washington County-Stakeholder Responses34

Table 26. Stakeholder Survey Results for Washington County and Maine
Table 27. Quantitative Health Indicators for Washington County, Maine and the U.S 41
Table 28. List of Data Sources and Years for Quantitative Health Indicators         47

# How to Use This Report

This report contains findings for Washington County from the Maine Shared Community Health Needs Assessment (Maine Shared CHNA) conducted in 2015. It is divided into ten sections to provide the reader with an easy-to-use reference to the data-rich assessment. It starts with the highest level of data, followed by summaries and synthesis of the data. The last sections include the detailed findings from assessments as well as the sources.

The report has several features that are important to keep in mind:

- The document provides a reference for more than 160 indicators and more than 30 qualitative survey questions covering many topics. It does not explore any individual topic in-depth.
- The definitions, sources and year(s) for each indicator discussed in the report are found at the end in the data sources section.
- Wherever the term, "statistically significant" is used to describe differences between data estimates, it means that the 95 percent confidence intervals for the given point estimates do not overlap.
- Unless otherwise noted, all rates presented in this report are age-adjusted and calculated per 100,000 population to facilitate comparisons between counties, Maine and the U.S.

The following is a brief description of each section.

#### Executive Summary

The summary provides the highest level overview of data for the county.

#### Background

This section explains the purpose and background of the SHNAPP and the Shared CHNA. It includes a description of the methodology and data sources used in the assessment.

#### County Demographics

The demographic section compares the population and socioeconomic characteristics of the county to the overall state of Maine.

#### Summary of Findings

This section provides a summary of the assessment data by health issue; it compares the county to the state and U.S. on key indicators and explains the importance of the health issues.

#### Stakeholder Feedback

High-level findings from the stakeholder survey are included in this section. It explores the top five health issues and factors identified as local priorities or concerns by stakeholders. It shares respondent concern for populations experiencing disparities in health status for these issues.

### Priority Health Issues and Challenges

Priority health issues and challenges appear in this section. This section categorizes the key findings from the quantitative and stakeholder (qualitative) datasets as strengths and challenges. The analysis includes health issue indicators from the quantitative datasets sorted into challenges and strengths, stakeholder responses for challenges and resources to address the challenges.

#### County Health Rankings

The 2015 County Health Ranking & Roadmaps model for the county is shown in this section. The model, from the University of Wisconsin Population Health Institute, shows how the individual health behaviors lead to health outcomes, which then determines the overall health status for a population. The graphic illustration includes the associated measures for each health indicator and the county rank among all 16 counties in the state of Maine. The data for the underlying health measures are those used by the University of Wisconsin in its 2015 report and may not always match the data shown in other sections of this report due to the time period for the data or use of different indicators.

#### Stakeholder Survey Findings

This section displays the full set of responses to each question asked in the stakeholder survey (excluding open-ended responses, which are available upon request). It compares the county to the statewide responses.

## Health Indicator Results from Secondary Data Sources

The results and sources section details the data for each of the 160 indicators for the county. It includes a table that compares data for the county, the state and the U.S. (where available). Statistically significant differences (at 95 percent confidence) are noted in this table where available and applicable.

#### Health Indicator Data Sources

This section lists the data source, year and additional notes for each indicator. In addition to the stakeholder survey conducted as a primary data source for this project, the secondary data sources used in this assessment include:

Child Maltreatment Report, Administration on Children Youth and Families
Maine Cancer Registry (MCR)
Maine Care
Maine Behavioral Risk Factor Surveillance System (BRFSS)
Maine CDC Drinking Water Program
Maine CDC HIV Program
Maine CDC Lead Program
Maine CDC National Electronic Disease Surveillance System (NEDSS)
Maine CDC Public Health Emergency Preparedness (PHEP)
Maine CDC STD Program

Maine CDC Vital Records Maine Department of Education Maine Department of Public Safety Maine Department of Labor Maine Department of Labor Maine Health Data Organization (MHDO) Maine Integrated Youth Health Survey (MIYHS) Maine Office of Data Research and Vital Records National Immunization Survey (NIS) National Survey of Children w/ Special Health Care Needs National Center for Health Statistics U.S. Bureau of Labor Statistics U.S. CDC WONDER & WISQARS U.S. Census

# **Executive Summary**

Public health and health care organizations share the goal of improving the lives of Maine people. Health organizations, along with business, government, community organizations, faith communities and individuals, have a responsibility to shape health improvement efforts based on sound data, personal or professional experience and community need.

This summary provides high-level findings from the Maine Shared Community Health Needs Assessment (CHNA), a comprehensive review of health data and community stakeholder input on a broad set of health issues in Maine. The Shared CHNA was conducted through a collaborative effort among Maine's four largest health-care systems – Central Maine HealthCare, Eastern Maine Healthcare Systems (EMHS), MaineGeneral Health, and MaineHealth – as well as the Maine Center for Disease Control and Prevention an office of the Maine Department of Health and Human Services (DHHS).

While it covers a broad range of topics, the Shared CHNA is not an exhaustive analysis of all available data on any single health issue. These data help identify priorities and should lead the reader to conduct a deeper investigation of the most pressing health issues.

Data are important and a solid starting point, but the numbers represent people who live in Maine. The overall goal of the Maine SHNAPP is to "turn data into action." Community engagement is therefore a critical next step, assuring shared ownership and commitment to collective action. The perspectives of those who live in our communities will bring these numbers to life and, together, we can set priorities to achieve measurable community health improvement. We invite all readers to use the information in this report as part of the solution to develop healthier communities in Maine.

## **Demographics and Socioeconomic Factors**

Washington County was home to 32,190 people in 2013. It is considered a rural county, according to the urban and rural classifications defined by the New England Rural Health RoundTable.<sup>1</sup> It is worse off than the state as a whole in many demographic and socioeconomic characteristics, including income, employment and poverty rates. Key demographic features include:

- Median household income of \$37,236 (2009-2013).
- 26.2 percent of children and 19.5 percent of all individuals live in poverty (2009-2013).
- The unemployment rate was 8.4 percent compared to the state at 5.7 percent (2014).

<sup>&</sup>lt;sup>1</sup> Rural Data for Action, New England Rural Health RoundTable, 2014. Available from: http://www.newenglandruralhealth.org/rural\_data

## Access to Health Care/Quality

Access to care in Washington County is worse that the state; specifically, a significantly lower percentage of Washington County residents have health insurance, although a lower proportion reported a lack of care due to cost. A higher proportion of adults and children have MaineCare as their medical insurance. The ambulatory care sensitive-conditions<sup>2</sup> hospital admission rate in Washington County was also significantly above the state. Key features for Washington County include:

- 13.7 percent of residents did not have health insurance (2009-2013); 9.6 percent experienced cost-related barriers to getting healthcare in the last year (2011-2013).
- 83.6 percent of adults reported having a personal doctor or other health care provider (2011-2013).
- 40.3 percent of adults were enrolled in MaineCare, 61.5 percent of children were enrolled (2015).
- The hospitalization rate for ambulatory care-sensitive conditions was 1,809 per 100,000 population (2011).

# General Health and Mortality

The general health of people in Washington County is worse than the state. Similar to the state, the top three leading causes of death in the county were cancer, heart disease and lower respiratory diseases. However, the county has a significantly higher overall mortality rate. In addition, a higher proportion of adults in Washington County report fair or poor health and more than one-third of adults have three or more chronic conditions. Key features for Washington County include:

- The overall mortality rate per 100,000 population was 875 in Washington County compared with 745.8 for the state (2009-2013).
- 20.4 percent of adults reported their health as fair or poor (2011-2013).
- 31.6 percent of adults have three or more chronic conditions (2011, 2013).

## Disease Incidence and Prevalence

Washington County has higher rates of chronic diseases incidence and prevalence than the state. Cancer is the leading cause of death in Washington County, with significantly higher rates of lung cancer but significantly lower rates of melanoma. Cardiovascular health is a concern in Washington County with significantly higher rates of heart attack mortality, coronary heart disease mortality and high cholesterol compared to the state. The county has higher rates of

<sup>&</sup>lt;sup>2</sup> Ambulatory care-sensitive conditions (ACSC) are Prevention Quality Indicators from the Agency for Healthcare Research and Quality and is intended to measure whether these conditions are being treated appropriately in the outpatient setting before hospitalization is required.

preventable diseases such as hepatitis B and C, and pertussis, but lower rates of chlamydia and Lyme disease than the state. Key features for Washington County include:

- Deaths due to any cancer were 210.5 per 100,000 population, compared to 185.5 statewide (2007-2011).
  - The number of new cases of all cancer sites per 100,000 population in Washington County was 533.3 (2007-2011). Lung cancer incidence was higher than the state, 92.8 versus 75.5 (2007-2011).
- For cardiovascular diseases:
  - The rate of acute myocardial infarction mortality per 100,000 population was 66.8 compared to 32.2 for the state (2009-2013).
  - The rate of coronary heart disease mortality was 130.8 per 100,000 population compared to the state at 89.8 (2009-2013).
  - The prevalence of high cholesterol was 45.9 percent compared to 40.3 percent for the state (2011, 2013).
- Newly reported cases of past or present hepatitis C per 100,000 population is 125.8 compared to 107.1 for the state (2014).
- Immunization exemptions among kindergarteners for philosophical reasons were 0.4 percent versus 3.7 percent statewide (2015).
- Pertussis incidence was 125.8 per 100,000 population in Washington County compared to 41.9 statewide (2014).
- Lyme disease incidence was 44 per 100,000 population (2014).

## Health Behaviors and Risk Factors

Overall, Washington County has slightly higher rates of people with mental health issues than the state. Alcohol and drug use among adults and high school students is similar to the state. Alcohol and drug induced mortality are also at higher rates as well as referrals for drug-affected babies. Current cigarette smoking rates among adults and high school students are slightly higher in Washington County than the state, while youth exposure to secondhand smoke is significantly higher in the county. The violent crime rate is higher than the state as a whole. Key health behavior and risk factors indicators for Washington County include:

- Mental health emergency department rates per 100,000 population were 2061.2 for the county compared to 1972.1 for the state (2011).
- Adult chronic heavy drinking was 8.3 percent (2011-2013). High school youth binge drinking was 17.8 percent (2013).
- Drug affected baby referrals received as a percentage of live births was 13.9 percent compared to 7.8 statewide (2014).
- Adult current cigarette smoking rates was 28.8 percent compared to 20.2 percent statewide (2011-2013).
- Youth secondhand smoke exposure was 53.2 percent compared to 38.3 percent statewide (2013).

• Washington County violent crime rate was 174 per 100,000 population compared to 125 statewide (2013).

# Stakeholder Priorities of Health Issues

Stakeholders who work in Washington County listed the following health issues as their top five concerns:

- Drug and alcohol abuse
- Obesity
- Tobacco use
- Mental health
- Cardiovascular diseases

Stakeholders identified the following populations as being disproportionately affected by the top health issues in Washington County:

- Low-income people, including those with incomes below the federal poverty level
- People with less than a high school education and/or low literacy (low reading or math skills)
- People who are medically underserved, including the uninsured and underinsured
- People with disabilities: physical, mental, or intellectual
- People in very rural and/or geographically isolated locations

Stakeholders prioritized the following factors as having a great influence on health in Washington County, resulting in poor health outcomes for residents:

- Poverty
- Employment
- Health care insurance
- Access to behavioral care/mental health care
- Health literacy

# Background

# Purpose

The Maine Shared Health Needs Assessment and Planning Process (SHNAPP) Project is a collaborative effort among Maine's four largest healthcare systems – Central Maine HealthCare, Eastern Maine Healthcare Systems (EMHS), MaineGeneral Health (MGH), and MaineHealth – as well as the Maine Center for Disease Control and Prevention (Maine CDC), an office of the Maine Department of Health and Human Services (Maine DHHS). The current collaboration expands upon the OneMaine Health Collaborative created in 2007 as a partnership among EMHS, MGH and MaineHealth. The Maine CDC and other partners joined these entities to develop a public-private partnership in 2012. The four hospital systems and the Maine CDC signed a memorandum of understanding in effect between June 2014 and December 2019 committing resources to the Maine SHNAPP Project.

The overall goal of the Maine SHNAPP is to "turn data into action" by conducting a shared community health improvement planning process for stakeholders across the state. The collaborative assessment and planning effort will ultimately lead to the implementation of comprehensive strategies for community health improvement. As part of the larger project, the Maine SHNAPP has pooled its resources to conduct this Shared Community Health Needs Assessment (Shared CHNA) to address community benefit reporting needs of hospitals, support state and local public health accreditation efforts, and provide valuable population health assessment data for use in prioritizing and planning for community health improvement.

This assessment builds on the earlier *OneMaine 2011 CHNA* that was developed by the University of New England and the University of Southern Maine, as well as the 2012 Maine State Health Assessment that was developed by the Maine DHHS. This Shared CHNA includes a large set of statistics on health status and risk factors from existing surveillance and health datasets. It differs from earlier assessments in two ways. Firstly, it includes input from a broad set of stakeholders from across the state from the 2015 SHNAPP Stakeholders' Survey. Secondly, it does not include the household telephone survey conducted for the OneMaine effort.

## Quantitative Data

This report contains both quantitative health data and qualitative stakeholder survey data on health issues and determinants affecting those living in Maine. The quantitative data come from numerous sources including surveillance surveys, inpatient and outpatient health data and disease registries. These data consist of 160 quantitative indicators within 18 groupings (domains) for reporting at the state level and, where possible, at the county and select urban levels. Please note that the data are taken from the most current year(s) available. Since the indicators come from a variety of sources, the data are measured over different time periods. In some cases, where there were not enough data in a single year to produce a statistically valid result, multiple years were combined to compute an indicator. Table 28 contains the complete list of the data sources.

## Qualitative Data

Qualitative data were collected through a statewide stakeholder survey conducted in May and June 2015 with 1,639 people representing more than 80 organizations and businesses in Maine. The survey was developed using a collaborative process that included Maine SHNAPP partners, Market Decisions Research and Hart Consulting, and a number of other stakeholders and health experts. In Washington County, a total of 133 stakeholders responded to the survey.

The objective of the survey was to produce qualitative data of the opinions of health professionals and community stakeholders on the health issues and needs of communities across the state. Given this purpose, the survey used a snowball sampling approach by inviting leaders of member organizations and agencies to invite their members and employees to participate. A concerted effort was made to recruit participants from a number of different industries and backgrounds across all communities in the state. Survey respondents represented public health and health care organizations as well as behavioral health, business, municipalities, education, public safety, and nongovernmental organizations. More than 80 organizations agreed to send the survey to their members or stakeholders.

The online survey was approximately 25 minutes in length and contained a number of questions about important health issues and determinants in the state, including a rating of most critical issues, the ability of Maine's health system (including public health) to respond to issues, availability of resources and assets to address specific health issues, impact on disparate populations, and identification of the entities primarily responsible for addressing issues and determinants. The survey asked all respondents a basic set of questions to rate the importance of health issues and impact of health factors. It then allowed respondents to provide answers to probing questions on the three issues and factors that they were most interested in or had the most knowledge about. Respondents provided over 12,000 open-ended comments to these indepth probing questions in the survey. The Market Decisions Research/Hart Consulting team reviewed, coded and cleaned all open-ended comments for similar and recurrent themes. Not all respondents shared comments for the probing questions.

## Limitations

While a number of precautions were taken to ensure that the results and findings presented in this report are sound and based upon statistically valid methods and analyses, there are some limitations to note. While the quantitative analysis used the most recent data sources available as of July 1, 2015, some of these sources contain data that are several years old. The most recent BRFSS and mortality data available at the time of analysis were from 2013, while the most recent hospitalization and cancer data were from 2011. This presents a particular challenge in trying to capture recent trends in health in the state, such as with opioid use. The data presented in this report may not necessarily represent the current situation in Maine, but are the best data available at the time of publication.

Given the qualitative nature of the survey questions and the sampling methodology, it is important to note that the results of the stakeholder survey are not necessarily representative of the population of Maine or a county at a given level of statistical precision. The findings reflect the informed opinions of health experts and community leaders from all areas of the state. However, it is important to use some caution when interpreting results, especially at the county level due to smaller sample sizes, as the results represent the opinions of only those who completed the survey.

# **Reports**

The Shared CHNA has several reports and datasets for public use that are available on the Maine CDC website and may be downloaded at <u>www.maine.gov/SHNAPP/</u>.


- <u>County-Level Maine Shared Community Health Needs Assessment Reports</u> summarize the data and provide insights into regional findings. These reports explore the priorities, challenges, and resources for each county and contain both summary and detailed tables.
- <u>State-Level Maine Shared Community Health Needs Assessment Report</u> includes information on each health issue, including analysis of sub-populations. The report includes state summaries and detailed tables.
- <u>Summary tables</u> are available for each public health district<sup>3</sup>, each county, and the cities of Portland and Bangor and the combined cities of Lewiston/Auburn.
- <u>Detailed Tables</u> contain each indicator, by subpopulation, region, and year.

<sup>&</sup>lt;sup>3</sup> To improve coordinated delivery of essential public health services, Department of Health and Human Services (DHHS) and the Maine Legislature approved the establishment of eight public health districts. District boundaries were established using population size, geographic areas, hospital service areas, and county borders. A District Liaison coordinates a Public Health Unit with co-located Maine CDC staff in one DHHS regional office for every District.


# **County Demographics**

Washington County has a total population of 32,190, with a slightly higher percentage of residents age 65 or more than the state of Maine as a whole. The county has the largest American Indian and Alaska Native population in the state, making up 5% of the total population. The socioeconomic and health characteristics of Washington County tend to be well below the state on many measures including income, poverty rates, unemployment and general health status.

# Figure 1. Population by Age Categories (U.S. Census 2013)<sup>4</sup>



# Figure 2. Population by Race/Ethnicity (U.S. Census 2013)



<sup>4</sup> Numbers may not add up to 100% due to rounding

# Washington County

Washington County is part of the Downeast Public Health District and Tribal Health District. Situated in eastern Maine, it hosts the easternmost point in the United States. The county is the third-least populous in Maine. Washington County is home to:

- Calais Regional Hospital.
- Down East Community Hospital.

Donulation	Washington		
Population	County	Maine	
Overall Population	32,190	1.33 mil	
Population density (per sq. mile)	12.8	43.1	
Percentage living in rural areas	100%	66.4%	
Single parent families	37.0%	34.0%	
65+ living alone	44.3%	41.2%	
Population living with a disability	20.5%	15.9%	
Economic Status			
Median household income	\$37,236	\$48,453	
Unemployment rate	8.4%	5.7%	
Adults and children living in poverty	19.5%	13.6%	
Children living in poverty	26.2%	18.5%	
Education			
HS graduation rate	89.4%	86.5%	

### **Key Demographics**

# Washington County Summary of Findings

# Socioeconomic Status

Economic opportunity and stability, including factors such income. as employment, food security and housing stability, have a significant impact on the health of individuals and communities. The Maine Behavioral 2013 Risk Factor Surveillance System (BRFSS) found the percentage of adults in Maine rating their health as excellent, very good or good was 94.8 percent among adults with household incomes of \$50,000 or more, but 53.8 percent among those with incomes under \$15,000.

In addition to income, there are many other social determinants of health, which have been defined as "conditions in the environments in which people are born, live, learn, work, play, worship and age that affect a wide range of health, functioning and quality-of-life outcomes and risks."<sup>5</sup> The conditions in which we live explain in part why some are healthier than others and why many generally are not as healthy as they could be. The Maine Shared CHNA takes into account a number of socioeconomic factors and other health determinants, including income and poverty, employment, education and household structure.

Percentage of adults and children living in poverty

Aroostool Franklin Hanco Oxford Kennebec York 9-11% 11-13% 13-15% 15-17% Maine Shared Health Needs 17% or more Assessment, 2015

<sup>&</sup>lt;sup>5</sup> The Institute of Medicine. Disparities in Health Care: Methods for Studying the Effects of Race, Ethnicity, and SES on Access, Use, and Quality of Health Care, 2002. Available from:

www.iom.edu/~/media/Files/Activity%20Files/Quality/NHDRGuidance/DisparitiesGornick.pdf

	Washington	Maine	U.S.
Adults and children living in poverty (2009-2013)	19.5%*	13.6%	15.4%
Children living in poverty (2009-2013)	26.2%*	18.5%	21.6%
Median household income (2009-2013)	\$37,236*	\$48,453	\$53 <i>,</i> 046
Single-parent families (2009-2013)	37.0%	34.0%	33.2%
65+ living alone (2009-2013)	44.3%	41.2%	37.7%

#### Table 1. Key Socioeconomic Indicators for Washington County

Asterisk (\*) and italics indicate a statistically significant difference between Washington County and Maine; NA = Not Available - data are not available for this indicator.

Note: U.S. results are from the most recently available year which may be different than county and state figures.

# General Health and Mortality

While it is essential to understand the causes, risk factors and other determinants of a population's health status, broad measures of health and mortality can also help explain the overall status and needs of the population in general and show in which populations there are disparities. General health status can be measured by self-reported data, as well as by mortality-related data such as life expectancy, leading causes of death and years of potential life lost.

	Washington	Maine	U.S.
Adults who rate their health fair to poor (2011-2013)	20.4%*	15.6%	16.7%
Adults with 14+ days lost due to poor mental health (2011-2013)	12.6%	12.4%	NA
Adults with 14+ days lost due to poor physical health (2011-2013)	14.4%	13.1%	NA
Adults with three or more chronic conditions (2011, 2013)	31.6%	27.6%	NA
Overall mortality rate per 100,000 population (2009-2013)	875.0*	745.8	731.9

#### Table 2. Key Health and Mortality Indicators for Washington County

Asterisk (\*) and italics indicate a statistically significant difference between Washington County and Maine; NA = Not Available - data are not available for this indicator.

Note: Age-adjusted rates presented in table; U.S. results are from the most recently available year which may be different than county and state figures.

The life expectancy in Washington County is 74.7 years for males and 80.1 years for females.

# Access to Health/Health Care Quality

Access to timely, appropriate, high-quality and regular health care and preventive health services is a key component of maintaining health. Good access to health care can be limited by financial, structural, and personal barriers. Access to health care is affected by location of and distance to health services, availability of transportation and the cost of obtaining the services – including the availability of insurance, the ability to understand and act upon information regarding

services, the cultural competency of health care providers and a host of other characteristics of the system and its clients. *Healthy People 2020* has identified four major components of access to health services: coverage, services, timeliness and workforce.<sup>6</sup>

In Washington County, 13.7 percent of residents did not have health insurance over the period from 2009-2013. However, access to health insurance does not necessarily guarantee access to care: among adults <u>with health insurance</u>, 5.9 percent in Washington County reported that they had experienced cost-related barriers to getting health care during the previous year (compared to 9.6 percent of all adults in the county).

	Washington	Maine	U.S.
Adults with a usual primary care provider (2011-2013)	83.6%*	87.7%	76.6%
Individuals who are unable to obtain or delay obtaining necessary medical care due to cost (2011-2013)	9.6%	11.0%	15.3%
Percent uninsured (2009-2013)	13.7%*	10.4%	11.7%
Ambulatory care-sensitive condition hospital admission rate per 100,000 population (2011)	1,809.0*	1,499.3	1,457.5
Adults with visits to a dentist in the past 12 months (2012)	56.9%*	65.3%	67.2%

Table 3. Key Access to Health/Health Care Quality Indicators for Washington County

Asterisk (\*) and italics indicate a statistically significant difference between Washington County and Maine; NA = Not Available - data are not available for this indicator.

Note: U.S. results are from the most recently available year which may be different than county and state figures.

Ambulatory care-sensitive hospital discharges is a Prevention Quality Indicator defined by the Agency for Healthcare Research and Quality (AHRQ) and is intended to measure whether conditions are being treated appropriately in the outpatient setting before hospitalization is required. AHRQ provides nationwide rates based on lower acuity and cost analysis of 44 states from the 2010 Agency for Healthcare Research and Quality's Healthcare Cost and Utilization Project State Inpatient Databases.<sup>7</sup>

# Chronic Disease

It is estimated that treatment for chronic diseases accounts for 86 percent of our nation's health care costs.<sup>8</sup> Chronic diseases include cancer, cardiovascular disease, diabetes and respiratory diseases like asthma and COPD, among other conditions. They are long-lasting health conditions and are responsible for seven out of ten deaths each year. Many chronic diseases can be

<sup>&</sup>lt;sup>6</sup> Healthy People 2020, Office of Disease Prevention and Health Promotion. Available from:

http://www.healthypeople.gov/2020/topics-objectives/topic/Access-to-Health-Services

<sup>&</sup>lt;sup>7</sup> Agency for Healthcare Research and Quality, Prevention Quality Indicators Technical Specifications - Version 5.0, March 2015, available at: http://www.qualityindicators.ahrq.gov/Modules/PQI\_TechSpec.aspx

<sup>&</sup>lt;sup>8</sup> National Center for Chronic Disease Prevention and Health Promotion, http://www.cdc.gov/chronicdisease/

prevented or controlled by reducing risk factors such as tobacco use, physical inactivity, poor nutrition and obesity.

Asthma is the most common childhood chronic condition in the United States and the leading chronic cause of children being absent from school.

Table 4. Key	Asthma and	COPD I	ndicators for	Washington Coun	tv
				, as a second second	·•.,

	Washington	Maine	U.S.
Asthma emergency department visits per 10,000 population (2009-2011)	118.0*	67.3	NA
COPD diagnosed (2011-2013)	8.6%	7.6%	6.5%
Current asthma (Adults) (2011-2013)	10.2%	11.7%	9.0%
Current asthma (Youth 0-17) (2011-2013)	10.9%	9.1%	NA
Pneumonia emergency department rate per 100,000 population (2011)	1,429.8*	719.9	NA
Pneumonia hospitalizations per 100,000 population (2011)	714.0*	329.4	NA

Asterisk (\*) and italics indicate a statistically significant difference between Washington County and Maine; NA = Not Available - data are not available for this indicator.

Note: Age-adjusted rates presented in table; U.S. results are from the most recently available year which may be different than county and state figures.

While the age-adjusted all-cancer incidence and mortality rates in Maine decreased significantly over the past ten years, cancer remains the leading cause of death among people in Maine. Cancer was also the leading cause of death in Washington County in 2013.

Table 5. Key	Cancer	Indicators	for `	Washington	County

	Washington	Maine	U.S.
Mortality – all cancers per 100,000 population (2007-2011)	210.5*	185.5	168.7
Incidence – all cancers per 100,000 population (2007-2011)	533.3	500.1	453.4
Mammograms females age 50+ in past two years (2012)	70.0%*	82.1%	77.0%
Colorectal screening (2012)	66.3%	72.2%	NA
Lung cancer incidence per 100,000 population (2007-2011)	92.8*	75.5	58.6
Melanoma incidence per 100,000 population (2007-2011)	11.1*	22.2	21.3

Asterisk (\*) and italics indicate a statistically significant difference between Washington County and Maine; NA = Not Available - data are not available for this indicator.

Note: Age-adjusted rates presented in table; U.S. results are from the most recently available year which may be different than county and state figures.

More than one in three adults lives with some type of cardiovascular disease. Heart disease and stroke can cause serious illness and disability with associated decreased quality of life and high economic costs. Cardiovascular disease conditions are among the most preventable health problems through the modification of common risk factors.

	Washington	Maine	U.S.
Acute myocardial infarction hospitalizations per 10,000 population (2010-2012)	39.4*	23.5	NA
Acute myocardial infarction mortality per 100,000 population (2009-2013)	66.8*	32.2	32.4
Cholesterol checked every five years (2011, 2013)	75.7%	81.0%	76.4%
Coronary heart disease mortality per 100,000 population (2009-2013)	130.8*	89.8	102.6
Heart failure hospitalizations per 10,000 population (2010-2012)	27.8*	21.9	NA
Hypertension prevalence (2011, 2013)	39.7%	32.8%	31.4%
High cholesterol (2011, 2013)	45.9%*	40.3%	38.4%
Stroke mortality per 100,000 population (2009-2013)	27.4	35.0	36.2

### Table 6. Key Cardiovascular Disease Indicators for Washington County

Asterisk (\*) and italics indicate a statistically significant difference between Washington County and Maine; NA = Not Available - data are not available for this indicator.

Note: Age-adjusted rates presented in table; U.S. results are from the most recently available year which may be different than county and state figures.

Diabetes mellitus is a complex health condition that lowers life expectancy, increases the risk of heart disease and is the leading cause of adult-onset blindness, lower-limb amputations and kidney failure. Lifestyle changes, effective self-management and treatment can delay or prevent diabetes and complications of diabetes.

#### **Table 7. Key Diabetes Indicators for Washington County**

	Washington	Maine	U.S.
Diabetes prevalence (ever been told) (2011-2013)	10.4%	9.6%	9.7%
Diabetes long-term complication hospitalizations (2011)	70.3	59.1	NA
Diabetes mortality (underlying cause) per 100,000 population (2009-2013)	33.9*	20.8	21.2

Asterisk (\*) and italics indicate a statistically significant difference between Washington County and Maine; NA = Not Available - data are not available for this indicator.

Note: Age-adjusted rates presented in table; U.S. results are from the most recently available year which may be different than county and state figures.

# **Environmental Health**

Environmental health includes the natural and built environments. Within these environments, there is risk of exposure to toxic substances and other physical hazards that exist in the air we breathe, the food we eat, the water we drink and the places where we live, play and work.<sup>9</sup>

Water quality issues in Maine include hazards such as disinfection byproducts, arsenic and nitrates/nitrites as well as bacteria contamination. Among households who get their drinking

<sup>&</sup>lt;sup>9</sup> Maine Center for Disease Control and Prevention. Healthy Maine 2020. Available from: http://www.maine.gov/dhhs/mecdc/healthy-maine/index.shtml

water from private wells, naturally occurring arsenic is a risk. Regular water quality testing can indicate the need for mitigation. In Washington County, 34.1 percent of households with private wells have tested their water for arsenic, compared with 43.3 percent of households statewide.

Childhood lead poisoning rates are of particular concern in areas with older housing. It can disproportionately affect people who live in older rental units and those who have less income.

	Washington	Maine	U.S.
Children with confirmed elevated blood lead levels (% among those screened) (2009-2013)	1.2%*	2.5%	NA
Children with unconfirmed elevated blood lead levels (% among those screened) (2009-2013)	5.3%*	4.2%	NA
Homes with private wells tested for arsenic (2009, 2012)	34.1%*	43.3%	NA
Lead screening among children age 12-23 months (2009- 2013)	79.5%*	49.2%	NA
Lead screening among children age 24-35 months (2009- 2013)	50.5%*	27.6%	NA

 Table 8. Key Environmental Health Indicators for Washington County

Asterisk (\*) and italics indicate a statistically significant difference between Washington County and Maine; NA = Not Available - data are not available for this indicator.

# **Immunization**

Immunization has accounted for significant decreases in morbidity and mortality of infectious diseases and an overall increase in life expectancy. However, many infectious diseases that can be prevented through vaccination continue to cause significant burdens on the health of Maine residents. The U.S. CDC has recommendations for a number of vaccines for young children, adolescents and older adults. Among its other recommendations, the U.S. CDC recommends yearly influenza vaccinations for people over six months of age.

	Washington	Maine	U.S.
Adults immunized annually for influenza (2011-2013)	42.0%	41.5%	NA
Adults immunized for pneumococcal pneumonia (ages 65 and older) (2011-2013)	68.0%	72.4%	69.5%
Immunization exemptions among kindergarteners for philosophical reasons (2015)	0.4%	3.7%	NA

Asterisk (\*) and italics indicate a statistically significant difference between Washington County and Maine; NA = Not Available - data are not available for this indicator.

Note: U.S. results are from the most recently available year which may be different than county and state figures.

# Infectious Disease/Sexually Transmitted Disease

There are 71 infectious diseases and conditions reportable in Maine. Surveillance data assist in monitoring trends in disease and identifying immediate threats to public health. However, there are limitations in surveillance data, specifically pertaining to underreporting. Available data reflects a subset of the disease burden in Maine.

Advances in sanitation, personal hygiene and immunizations have provided control over some diseases, but others continue to thrive despite best efforts. Lyme disease, if left untreated, can cause severe headaches, severe joint pain and swelling, inflammation of the brain and short-term memory problems<sup>10</sup>. Incidence has increased from 224 reported cases statewide in 2004 to 1,400 in 2014, a growth of more than 500 percent in a decade.

#### Table 10. Key Infectious Disease Indicators for Washington County

	Washington	Maine	U.S.
Incidence of past or present hepatitis C virus (HCV) per 100,000 population (2014)	125.8	107.1	NA
Incidence of newly reported chronic hepatitis B virus (HBV) per 100,000 population (2014)	0.0	8.1	NA
Lyme disease incidence per 100,000 population (2014)	44.0	105.3	10.5

Asterisk (\*) and italics indicate a statistically significant difference between Washington County and Maine; NA = Not Available - data are not available for this indicator.

Note: U.S. results are from the most recently available year which may be different than county and state figures.

While the rates of sexually transmitted diseases like chlamydia, gonorrhea and HIV are significantly lower in Maine than the U.S., it is an issue that disproportionately affects specific segments of the population, including young adults and men who have sex with men.

Table 11	. Kev	Sexually	<b>Transmitted</b>	l Disease	<b>Indicators</b>	for	Washington	County
				~ ~ ~ ~ ~ ~ ~				

	Washington	Maine	U.S.
Chlamydia incidence per 100,000 population (2014)	157.2	265.5	452.2
Gonorrhea incidence per 100,000 population (2014)	15.7	17.8	109.8
HIV incidence per 100,000 population (2014)	0.0	4.4	11.2

Asterisk (\*) and italics indicate a statistically significant difference between Washington County and Maine; NA = Not Available - data are not available for this indicator.

Note: U.S. results are from the most recently available year which may be different than county and state figures.

# Injuries

Intentional or violence-related injury is an important public health problem that affects people of all ages. Violence prevention activities include changing societal norms regarding the

<sup>&</sup>lt;sup>10</sup> Signs and Symptoms of Untreated Lyme Disease, Centers for Disease Control and Prevention (CDC), Available from: http://www.cdc.gov/lyme/signs\_symptoms/

acceptability of violence, improving conflict resolution and other problem-solving skills and developing policies to address economic and social conditions that can lead to violence.

Suicide is the second leading cause of death among 15- to 34-year-olds in Maine and the tenth leading cause of death among all ages combined. In Washington County, the age-adjusted rate of suicide deaths was 21.1 per 100,000 population, compared to 15.2 for the state over the same time period.

	Washington	Maine	U.S.		
Domestic assault reports to police per 100,000 population (2013)	394.6	413.0	NA		
Firearm deaths per 100,000 population (2009-2013)	17.0*	9.2	10.4		
Suicide deaths per 100,000 population (2009-2013)	21.1	15.2	12.6		
Violent crime rate per 100,000 population (2013)	174.0	125.0	367.9		
		<i>a</i> , 1	17 .		

Table 12. Key	Intentional I	niurv I	Indicators	for W	Vashington	County
1 abic 12. Key	intentional i	injuryi	inuicators	IUI V	asinington	County

Asterisk (\*) and italics indicate a statistically significant difference between Washington County and Maine; NA = Not Available - data are not available for this indicator.

Note: Age-adjusted rates presented in table; U.S. results are from the most recently available year which may be different than county and state figures.

Unintentional injuries are a leading cause of death and disability. While many people think of unintentional injuries as a result of accidents, most are predictable and preventable. Unintentional injury was the leading cause of death among 1- to 44-year-olds in Maine and the fifth-leading cause of death among all ages combined in 2013. Motor vehicle crashes, unintentional poisonings, traumatic brain injuries and falls lead to millions of dollars in medical and lost work costs.

	Washington	Maine	U.S.
Always wear seatbelt (Adults) (2013)	75.8%*	85.2%	NA
Always wear seatbelt (High School Students) (2013)	42.6%*	61.6%	54.7%
Traumatic brain injury related emergency department visits (all intents) per 10,000 population (2011)	67.9*	81.4	NA
Unintentional and undetermined intent poisoning deaths per 100,000 population (2009-2013)	15.6	11.1	13.2
Unintentional fall related injury emergency department visits per 10,000 population (2011)	415.6*	361.3	NA
Unintentional motor vehicle traffic crash related deaths per 100,000 population (2009-2013)	19.5*	10.8	10.5

## Table 13. Key Unintentional Injury Indicators for Washington County

Asterisk (\*) and italics indicate a statistically significant difference between Washington County and Maine; NA = Not Available - data are not available for this indicator.


Note: Age-adjusted rates presented in table; U.S. results are from the most recently available year which may be different than county and state figures.

# Mental Health

Mental health is a complex issue that can affect many facets of a person's daily life. In the U.S., about one in four adults and one in five children have diagnosable mental disorders and they are the leading cause of disability among people ages 15-44.<sup>11</sup> In Washington County, 20.1 percent of adults reported currently receiving outpatient mental health treatment (taking medicine or receiving treatment from a doctor) in 2011-2013, compared to 17.7 percent of adults statewide.

Mental well-being can also affect a person's physical health in many ways, including chronic pain, a weakened immune system and increased risk for cardiovascular problems. In addition, mental illnesses, such as depression and anxiety, affect people's ability to participate in health-promoting behaviors.<sup>12</sup>

Stigma, additional health issues, access to services and complexities of treatment delivery also prevent many from receiving adequate treatment for their mental health issues.



Percentage of Adults with Current Depression

Table 14. Key Mental Health Indicators for Washington (	County
---	--------

	Washington	Maine	U.S.
Adults who have ever had depression (2011-2013)	23.0%	23.5%	18.7%
Adults with current symptoms of depression (2011-2013)	11.4%	10.0%	NA
Adults currently receiving outpatient mental health treatment (2011-2013)	20.1%	17.7%	NA
Sad/hopeless for two weeks in a row (High School Students) (2013)	28.7%	24.3%	29.9%
Seriously considered suicide (High School Students) (2013)	15.4%	14.6%	17.0%

Asterisk (\*) and italics indicate a statistically significant difference between Washington County and Maine; NA = Not Available - data are not available for this indicator.

Note: Age-adjusted rates presented in table; U.S. results are from the most recently available year which may be different than county and state figures.

<sup>&</sup>lt;sup>11</sup> Guide to Community Preventive Services. Improving mental health and addressing mental illness. www.thecommunityguide.org/mentalhealth/index.html.

<sup>&</sup>lt;sup>12</sup> US Department of Health and Human Services. Health People 2020: Mental Health and Mental Disorders. 2012 Available from: www.healthypeople.gov/2020/topicsobjectives2020/overview.aspx?topicid=28.

# Physical Activity, Nutrition and Weight

Eating a healthy diet, being physically active and maintaining a healthy weight are essential for an individual's overall health. These three factors can help lower the risk of developing numerous health conditions, including high cholesterol, high blood pressure, heart disease, stroke, diabetes and cancer. They also can help prevent existing health conditions from worsening over time.

Sugar-sweetened beverages, such as non-diet soda, sports drinks and energy drinks, provide little to no nutritional value, but their calories can lead to obesity and being overweight, along with health risks including tooth decay, heart disease and type 2 diabetes

The 2008 *Physical Activity Guidelines for Americans* recommends that adults, age 18-64, get a minimum of 150 minutes of moderateintensity physical activity a week and that children, age 6-17, get 60 or more minutes of physical activity each day.<sup>13</sup> Among adults in Washington County from 2011-2013, 27.9 percent led a sedentary lifestyle, meaning they did not participate in any leisure-time (nonwork) physical activity or exercise in the previous month.




Table 15. Key	v Nutrition and	<b>Physical Activ</b>	vity Indicators fo	or Washington County
	,			

	Washington	Maine	U.S.
Fruit and vegetable consumption (High School Students) (2013)	14.0%	16.8%	NA
Fruit consumption among Adults 18+ (less than one serving per day) (2013)	35.4%	34.0%	39.2%
Met physical activity recommendations (Adults) (2013)	45.9%	53.4%	50.8%
Physical activity for at least 60 minutes per day on five of the past seven days (High School Students) (2013)	41.6%	43.7%	47.3%

<sup>13</sup> Physical Activity Guidelines for Americans, U.S. Department of Health and Human Services, 2008, http://health.gov/Paguidelines/guidelines/

	Washington	Maine	U.S.
Sedentary lifestyle – no leisure-time physical activity in past month (Adults) (2011-2013)	27.9%*	22.4%	25.3%
Soda/sports drink consumption (High School Students) (2013)	34.7%*	26.2%	27.0%
Vegetable consumption among Adults 18+ (less than one serving per day) (2013)	19.7%	17.9%	22.9%

Asterisk (\*) and italics indicate a statistically significant difference between Washington County and Maine; NA = Not Available - data are not available for this indicator.

Note: U.S. results are from the most recently available year which may be different than county and state figures.

In 2013, 65.4 percent of adults 18 years and older in Washington County were overweight or obese (35.0 percent were overweight and 30.4 percent were obese). Overall in Maine, 64.8 percent of adults were overweight or obese.

#### Table 16. Key Weight Indicators for Washington County

	Washington	Maine	U.S.
Obesity (Adults) (2013)	30.4%	28.9%	29.4%
Obesity (High School Students) (2013)	15.9%	12.7%	13.7%
		~ .	

Asterisk (\*) and italics indicate a statistically significant difference between Washington County and Maine; NA = Not Available - data are not available for this indicator.

Note: U.S. results are from the most recently available year which may be different than county and state figures.

# **Pregnancy and Birth Outcomes**

Addressing health risks during a woman's pregnancy can help prevent future health issues for women and their children. Increasing access to quality care both before pregnancy and between pregnancies can reduce the risk of pregnancy-related complications and maternal and infant mortality. Early identification and treatment of health issues among babies can help prevent disability or death.<sup>14</sup>

#### Table 17. Key Pregnancy and Birth Outcomes for Washington County

	Washington	Maine	U.S.
Infant deaths per 1,000 live births (2003-2012)	4.3	6.0	6.0
Live births for which the mother received early and adequate prenatal care (2010-2012)	84.3%	86.4%	84.8%
Live births to 15-19 year olds per 1,000 population (2010-2012)	28.7*	20.5	26.5
Low birth weight (<2500 grams) (2010-2012)	4.7%	6.6%	8.0%

Asterisk (\*) and italics indicate a statistically significant difference between Washington County and Maine; NA = Not Available - data are not available for this indicator.

Note: U.S. results are from the most recently available year which may be different than county and state figures.

<sup>14</sup> Healthy People 2020. Maternal, infant, and child health: overview. Available from:

http://www.healthypeople.gov/2020/topics-objectives/topic/maternal-infant-and-child-health


# Substance and Alcohol Abuse

Substance abuse and dependence are preventable health risks that lead to increased medical costs, injuries, related diseases, cancer and even death. Substance abuse also adversely affects productivity and increases rates of crime and violence.<sup>15</sup> In Maine in 2010, approximately \$300 million was spent on medical care where substance use was a factor.<sup>16</sup>

Of particular note is the recent increase in heroin and prescription opioid dependence and mortality, both nationally and in the state. From 2002 to 2013, heroin overdose death rates nearly quadrupled in the U.S., from 0.7 deaths to 2.7 deaths per 100,000 population. The rates nearly doubled from 2011 to 2013.<sup>17</sup> In addition, data from the National Survey on Drug Use and Health (NSDUH) indicate that heroin use, abuse and dependence have increased in recent years.<sup>11</sup>

The heroin problem in Maine has become a focus of national attention.<sup>18</sup> Deaths from heroin overdoses in Maine rose from seven in 2010 to 57 in 2014<sup>19</sup> and that number continues to climb in 2015.<sup>20</sup>

**Substance Abuse Hospitalizations** 



<sup>&</sup>lt;sup>15</sup> National Institute on Drug Abuse. Principles of Drug Abuse Treatment for Criminal Justice Populations: A Research-Based Guide. Bethesda, MD: National Institutes of Health, National Institute on Drug Abuse. NIH publication No. 11-5316, revised 2012. Available at www.drugabuse.gov/publications/principles-drug-abuse-treatment-criminal-justice-populations

<sup>&</sup>lt;sup>16</sup> The Cost of Alcohol and Drug Abuse in Maine, 2010. Office of Substance Abuse and Mental Health Services, Department of Health and Human Services, 2013. Available at: http://www.maine.gov/dhhs/samhs/osa/pubs/data/2013/Cost2010-final%20Apr%2010%2013.pdf

<sup>&</sup>lt;sup>17</sup> Jones CM, Logan J, Gladden M, Vital Signs: Demographic and Substance Use Trends Among Heroin Users — United States, 2002–2013, Morbidity and Mortality Weekly Report (MMWR), 2015. Available from:

http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6426a3.htm

<sup>&</sup>lt;sup>18</sup> Heroin in New England, More Abundant and Deadly. The New York Times. July 18, 2013.

http://www.nytimes.com/2013/07/19/us/heroin-in-new-england-more-abundant-and-deadly.html

<sup>&</sup>lt;sup>19</sup> Heroin Deaths in Maine Jump – Record Level of Overdose Deaths in 2014. May 15, 2015. Office of the Chief Medical Examiner (OCME) of the Office of the Maine Attorney General. Available at:

http://www.maine.gov/ag/news/article.shtml?id=644190

<sup>&</sup>lt;sup>20</sup> First half of 2015 shows pace of drug deaths has not slowed – Heroin, Fentanyl deaths continue to surge. August 20, 2015. Office of the Chief Medical Examiner (OCME) of the Office of the Maine Attorney General. Available at:

http://www.maine.gov/ag/news/article.shtml?id=653671

	Washington	Maine	U.S.
Alcohol-induced mortality per 100,000 population (2009-2013)	11.0	8.0	8.2
Chronic heavy drinking (Adults) (2011-2013)	8.3%	7.3%	6.2%
Drug-affected baby referrals received as a percentage of all live births (2014)	13.9%	7.8%	NA
Drug-induced mortality per 100,000 population (2009-2013)	18.7	12.4	14.6
Emergency medical service overdose response per 100,000 population (2014)	389.8	391.5	NA
Opiate poisoning (ED visits) per 100,000 population (2009-2011)	20.8	25.1	NA
Past-30-day alcohol use (High School Students) (2013)	28.2%	26.0%	34.9%
Past-30-day marijuana use (High School Students) (2013)	23.6%	21.6%	23.4%
Prescription Monitoring Program opioid prescriptions (days supply/pop) (2014-2015)	9.0	6.8	NA
Substance-abuse hospital admissions per 100,000 population (2011)	360.0	328.1	NA

#### Table 18. Key Substance Abuse Indicators for Washington County

Asterisk (\*) and italics indicate a statistically significant difference between Washington County and Maine; NA = Not Available - data are not available for this indicator.

Note: Age-adjusted rates presented in table; U.S. results are from the most recently available year which may be different than county and state figures.

# Tobacco Use

Use of tobacco is the most preventable cause of disease, death and disability in the United States. Despite this, more than 480,000 deaths in the United States are attributable to tobacco use every year <sup>21</sup> (more than from alcohol use, illegal drug use, HIV, motor vehicle injuries and suicides combined). In addition, exposure to secondhand tobacco smoke has been causally linked to cancer and to respiratory and cardiovascular diseases in adults, and to adverse effects on the health of infants and children, such as respiratory and ear infections.<sup>22</sup> While the percentage of Maine adults who smoke cigarettes has declined significantly over time, one-fifth of the state's population still smokes cigarettes, including 28.8 percent of adults in Washington County.

<sup>&</sup>lt;sup>21</sup> U.S. Department of Health and Human Services. The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General. Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2014

<sup>&</sup>lt;sup>22</sup> U.S. Department of Health and Human Services. Healthy People 2020.

Leading health indicators: tobacco overview and impact. Available from:

http://www.healthypeople.gov/2020/LHI/tobacco.aspx

	Washington	Maine	U.S.
Current smoking (Adults) (2011-2013)	28.8%	20.2%	19.0%
Current smoking (High School Students) (2013)	19.7%	12.9%	15.7%
Current tobacco use (High School Students) (2013)	23.6%	18.2%	22.4%
Secondhand smoke exposure (Youth) (2013)	53.2%*	38.3%	NA

### Table 19. Key Tobacco Use Indicators for Washington County

Asterisk (\*) and italics indicate a statistically significant difference between Washington County and Maine; NA = Not Available - data are not available for this indicator.

Note: U.S. results are from the most recently available year which may be different than county and state figures.

# Stakeholder Feedback

In June 2015, the Maine Shared CHNA research team conducted a survey among stakeholders to identify and prioritize significant health issues in communities across the state. The purpose of the survey was to include the voices and broad interests of local stakeholders about community health needs in their areas. The survey instrument was designed in collaboration with the Maine Shared CHNA Steering Committee and its work groups; it covered four domains of questions:

- Stakeholder demographic information
- Health issues with the greatest impact
- Determinants of health
- Health priorities and challenges

The survey was administered using a snowball approach, where stakeholder agencies agreed to send the surveys to their members and stakeholders for participation. Statewide, 1,639 people completed the survey; 133 of the total respondents indicated that they worked in Washington County or the Downeast Public Health District. Respondents represented health care agencies, public health agencies, law enforcement, municipalities, schools, businesses, social service agencies and nongovernmental organizations.

There were 403 respondents who indicated they worked at the state-level (e.g., Maine CDC, businesses that spanned the state, etc.). These respondents were included in the overall results, but were not included in any of the county-level results. Respondents could indicate that they represent more than one county in the survey, therefore the total of completed surveys by county will add up to more than 1,639.

## **Stakeholder Ratings of Health Issues**

How much of a problem is \_\_\_\_ in Washington County? (Responses were provided on a 5 point scale where 1-Not at all a problem, 2-Minor problem, 3-Moderate problem, 4-Major problem, 5-Critical problem (This table includes % reporting 4-Major or 5-Critical problem).

Health Issue	Washington	Maine
Family Health	n=133	n=1639
Elder health	64%	55%
Childhood obesity	49%	58%
Child developmental issues	44%	34%
Maternal and child health	31%	23%
Adolescent health	32%	25%
Infant mortality	6%	4%
Chronic Diseases	1	<u> </u>
Obesity	82%	78%
Cardiovascular diseases	76%	63%
Depression	74%	67%
Diabetes	74%	63%
Cancer	74%	50%
Respiratory diseases	69%	60%
Musculoskeletal diseases	40%	28%
Neurological diseases	38% 355	
Infectious Diseases		
Infectious diseases	25%	22%
Sexually transmitted		
diseases/HIV/AIDS	14%	13%
Healthy Behaviors		
Drug and alcohol abuse	91%	80%
Tobacco use	77%	63%
Physical activity and		
nutrition	74%	69%
Other Health Issues		
Mental health	76%	71%
Oral health	67%	53%
Unintentional injury	48%	34%
Suicide and self-harm	46%	37%
Violence	42%	38%
Lead poisoning and other		
environmental health issues	23%	17%

# **Top Health Issues**


Washington County stakeholders ranked a set of 25 health issues on "how you feel they impact overall health of residents" on a five-point scale, where 1 is "not at all a problem" and 5 is "critical problem." The top five issues of concern reported for the county were:


- Drug and alcohol abuse
- Obesity
- Tobacco use
- Mental health
- Cardiovascular diseases

Respondents were asked probing statements about the three issues they knew the most about. The question was worded as follows:

"The health system (including public health) in Washington County has the ability to significantly improve [] health issue."

Stakeholder responses on the probing question for the top five health issues appear in Figure 3.





Maine Shared Community Health Needs Assessment, 2015

\* Results presented for Maine due to small sample size at the county level

Stakeholders were also asked to share their thoughts on the populations experiencing health disparities for the health issues that they selected. Results for the top five health issues in Washington County are presented in Table 20.

Table 20. Percentage of Stakeholders who agreed that Significant Disparities Exist Among
Specific Groups for a Specific Health Issue.

Populations Experiencing Health Disparities	Cardiovascular diseases	Obesity	Drug and alcohol abuse	Tobacco use	Mental health
Low- income, including those below the federal poverty limit	80%	87%	85%	90%	79%
Medically-underserved - including uninsured and under-insured	78%	70%	63%	62%	74%
Less than a high school education and/ or low literacy	52%	61%	67%	74%	56%
Very rural and/or geographically isolated people	54%	44%	49%	40%	56%
People with disabilities - physical, mental, or intellectual	37%	47%	41%	37%	63%
Limited or no English proficiency	21%	12%	14%	16%	21%
Military veterans	14%	4%	34%	29%	43%
Gay, lesbian, bisexual or transgendered people	3%	4%	30%	33%	36%
Women	18%	15%	17%	8%	20%
Members of any Federally-recognized Tribe	11%	12%	21%	15%	19%
Refugees/immigrants	10%	4%	8%	3%	20%
Specific age group	15%	10%	12%	9%	12%
Racial/ethnic minority populations	8%	4%	9%	1%	11%
Deaf and hard of hearing people	3%	3%	3%	-	11%
Adolescents/Teens (13-17)	1%	3%	8%	3%	6%
Seniors/Elderly (65+)	10%	3%	-	2%	3%
Youth/Children (0-12)	-	4%	-	1%	4%
Adults (21-64)	6%	1%	3%	2%	-
Young adults (18-21)	1%	1%	7%	3%	2%
Other	6%	6%	12%	9%	12%

Stakeholder input also pointed out the key social or environmental drivers in Maine that lead to these health issues. The key drivers for the top five health issues in Washington County are presented in Table 21.

icua to a specific ficardi condition					
Key Drivers	Cardiovascular diseases	Obesity	Drug and alcohol abuse	Tobacco use	Mental health
Poverty/low income/low socio-economic status	36%	40%	30%	40%	27%
Lack of education	25%	31%	11%	43%	15%
Lack of access to healthy foods	15%	28%	-	-	1%
Bad eating habits	26%	26%	-	-	1%
Lack of access to physical activity opportunities	7%	25%	-	-	-
Lack of access to behavioral care/mental health care	1%	-	3%	1%	44%
Isolated and rural areas	7%	9%	11%	6%	14%
Inadequate health literacy	12%	9%	8%	7%	-
Cultural or social norms/acceptance/role modeling	5%	9%	22%	38%	4%
Lack of transportation	12%	8%	6%	3%	11%
Lack of access to treatment	7%	2%	33%	7%	2%
Lack of employment opportunities	2%	2%	17%	7%	6%
Social attitudes such as discrimination, stigma, etc.	2%	2%	14%	7%	34%
Lack of health care insurance	9%	2%	5%	1%	10%
Adverse childhood experiences	1%	2%	3%	1%	5%
Substance use/addiction	22%	2%	2%	10%	5%
Lack of access to primary care	22%	2%	-	3%	3%
Personal responsibility	5%	8%	4%	1%	3%
Apathy/depression/hopelessness	2%	5%	11%	6%	2%
Food insecurity	-	4%	-	1%	1%
Co-morbidity-physical or behavioral	7%	3%	-	3%	4%
Lack of exercise	12%	3%	-	-	-
Lack of social support and interactions-positive	-	2%	14%	1%	1%
Mental illness	2%	2%	2%	6%	2%
Lack of civic participation	-	2%	-	-	1%
Abuse/trauma	-	1%	3%	-	3%
Lack of funding-programs/low reimbursement to providers	-	1%	2%	3%	8%

Table 21. Percentage of Stakeholders who identified Certain Factors as Key Drivers that
lead to a Specific Health Condition

The next section of this report has a list of the community resources and assets that are available in the area to address these health issues and drivers, along with a summary of the additional resources that are needed. See **Table 23. Priority Health Issues** in the following section of this report.

# **Top Health Factors**

Health factors are those conditions, such as health behaviors, socioeconomic status, or physical environment features that can affect the health of individuals and communities. Stakeholders prioritized 26 health factors in five categories that can play a significant role in the incidence and prevalence of health problems in their communities.

Stakeholders responded to the following question: "For the factors listed below, please indicate how much of a problem each is in your area and leads to poor health outcomes for residents." They responded using a scale of 1 to 5, where 1 means "not a problem at all," and 5 means "critical problem." Respondents selected the following five factors as greatest problems that lead to poor health outcomes in Washington County:

- Poverty
- Employment
- Health care insurance
- Access to behavioral care/mental health care
- Health literacy

As with health issues, stakeholders were asked further probing questions on the three factors that they believe have the greatest impact on the health of their county.

To understand the capacity available in the county to address the most significant health factors identified by stakeholders, respondents were asked additional probing statements about the issues they knew the most about. "The health system (including public health) in Washington County has the ability to significantly improve these health factors with the current investment of time and resources."

## **Stakeholder Ratings of Health Factors**

How much of a problem is \_\_ in Washington County? (Responses were provided on a 5 point scale where 1-Not at all a problem, 2-Minor problem, 3-Moderate problem, 4-Major problem, 5-Critical problem (This table includes % reporting 4-Major or 5-Critical problem).

problem).		
Health Factor	Washington	Maine
Economic Stability	n=133	n=1639
Poverty	89%	78%
Employment	82%	64%
Food security	66%	58%
Housing stability	59%	57%
Education		
Early Childhood		
Education/Development	57%	43%
Enrollment in higher		
education	41%	35%
Language and literacy	39%	34%
High school graduation	35%	31%
Social and Community Contex	ſt	
Adverse childhood		
experiences	64%	56%
Social support and		
interactions	53%	50%
Caregiver support	48%	46%
Social Attitudes (such as		
Discrimination)	46%	38%
Incarceration or		
Institutionalization	35%	35%
Civic participation	34%	30%
Health and Health Care		
Health care insurance	78%	64%
Access to behavioral		
care/mental health care	76%	67%
Health literacy	74%	62%
Access to oral health	66%	56%
Access to other health care	57%	41%
Access to primary care	56%	39%
Neighborhood and Built Environment		
Transportation	70%	67%
Access to physical activity		
opportunities	59%	42%
Access to healthy foods	58%	53%
Quality of housing	38%	34%
Crime and violence	37%	27%
Environmental Conditions		
(Air quality, water quality,		
pollution, etc.)	16%	12%
· · ·	1	

Stakeholder responses on the probing question for the top five health issues appear in Figure 4.




Figure 4. The health system (including public health) has the ability to significantly improve these health factors with the current investment of time and resources.

Maine Shared Community Health Needs Assessment, 2015

\* Results presented for Maine due to small sample size at the county level

The next section of this report has a list of the community resources and assets that are available in the area to address these health factors, along with a summary of the additional resources that are needed. See **Table 25. Priority Health Factors** in the next section.

## Washington County Priority Health Issues and Factors

Table 22 presents a summary of the health issues - successes and challenges - experienced by residents of Washington County. Data come from a comprehensive analysis of available surveillance data (see Table 28 for a full list of the health indicators and factors included in this project). Two criteria were used to select the issues in this table: statistically significant and relative differences between the county and state. **Statistically significant differences**, using a 95 percent confidence level, are noted with an asterisk (\*) after the indicator. A **rate ratio** was also calculated to compare the relative difference between the county and state. Indicators where the county was 15 percent or more above or below the state average are included in this table.

 
 Table 22. Priority Health Issue Successes and Challenges for Washington County-Surveillance Data

Health Issues - Surveillance Data					
Health Successes	Health Challenges				
• Low incidence of bladder cancer [WAS=24.0; ME=28.3] and melanoma per 100,000 population [WAS=11.1; ME=22.2]*	Compared to the state, Washington County faces numerous health challenges across many health areas. Some of these challenges include:				
• Low stroke mortality per 100,000 population [WAS=27.4; ME=35.0]	<ul> <li>Higher percentage of adults who rate their health fair to poor [WAS=20.4%; ME=15.6%]*</li> </ul>				
• Fewer children with confirmed elevated blood lead levels (% among those screened)	<ul> <li>High overall mortality rate per 100,000 population [WAS=875.0; ME=745.8]*</li> </ul>				
<ul> <li>[WAS=1.2%; ME=2.5%]*</li> <li>Lower incidence of some infectious and sexually transmitted diseases, including</li> </ul>	<ul> <li>High ambulatory care-sensitive condition hospital admission rate per 100,000 population [WAS=1,809.0; ME=1,499.3]*</li> </ul>				
<ul> <li>Newly reported chronic hepatitis B virus (HBV) [WAS=0.0; ME=8.1] and Lyme disease per 100,000 population [WAS=44.0; ME=105.3]</li> </ul>	<ul> <li>A number respiratory issues in adults and children, including more asthma emergency department visits per 10,000 population [WAS=118.0; ME=67.3]*, higher percentage of current asthma among children</li> </ul>				
<ul> <li>Chlamydia [WAS=157.2; ME=265.5] and HIV incidence per 100,000 population [WAS=0.0; ME=4.4]</li> </ul>	<ul> <li>(0-17) [WAS=10.9%; ME=9.1%], as well as a high pneumonia emergency department rate</li> <li>[WAS=1,429.8; ME=719.9]* and high pneumonia hospitalizations per 100,000 population [WAS=714.0;</li> </ul>				
• Lower rates of reported rape per 100,000 population [WAS=9.3; ME=27.0]	<ul><li>ME=329.4]*</li><li>High mortality for a number of different cancers,</li></ul>				
• Fewer traumatic brain injury related emergency department visits (all intents) per 10,000 population [WAS=67.9; ME=81.4]*	<ul> <li>All cancers per 100,000 population [WAS=210.5; ME=185.5]*</li> </ul>				
• Low rates of infant deaths per 1,000 live births [WAS=4.3; ME=6.0]	<ul> <li>Female breast cancer per 100,000 population [WAS=23.8; ME=20.0]</li> </ul>				

Health Issues - Surveillance Data						
Health Successes Health Challenges						
<ul> <li>Lower low birth weight (&lt;2500 grams)</li> </ul>	<ul> <li>Prostate cancer per 100,000 population [WAS=29.1; ME=22.1]</li> </ul>					
<ul> <li>[WAS=4.7%; ME=6.6%]</li> <li>Low opiate poisoning ED visits [WAS=20.8; ME=25.1] and hospitalizations per 100,000 population [WAS=11.1; ME=13.2]</li> </ul>	<ul> <li>Tobacco-related neoplasms per 100,000 population [WAS=46.3; ME=37.4]</li> </ul>					
	<ul> <li>High incidence of various cancers, including:</li> <li>Colorectal late-stage incidence per 100,000 population [WAS=26.4; ME=22.7]</li> </ul>					
• Lower percentage of past-30-day nonmedical use of prescription drugs among high school students [WAS=4.2%; ME=5.6%]	<ul> <li>Colorectal cancer incidence per 100,000 population [WAS=52.8; ME=43.5]</li> </ul>					
	<ul> <li>Lung cancer incidence per 100,000 population [WAS=92.8; ME=75.5]*</li> </ul>					
	<ul> <li>High levels of cardiovascular risk factors and diseases, which includes:</li> </ul>					
	<ul> <li>Acute myocardial infarction hospitalizations per 10,000 population [WAS=39.4; ME=23.5]*</li> </ul>					
	<ul> <li>Acute myocardial infarction mortality per 100,000 population [WAS=66.8; ME=32.2]*</li> </ul>					
	<ul> <li>Coronary heart disease mortality per 100,000 population [WAS=130.8; ME=89.8]*</li> </ul>					
	<ul> <li>Heart failure hospitalizations per 10,000 population [WAS=27.8; ME=21.9]*</li> </ul>					
	<ul> <li>Higher levels of hypertension prevalence [WAS=39.7%; ME=32.8%]</li> </ul>					
	<ul> <li>Higher percentage of adults with high cholesterol [WAS=45.9%; ME=40.3%]*</li> </ul>					
	<ul> <li>Higher rates of diabetes emergency department visits (principal diagnosis) per 100,000 population [WAS=294.1; ME=235.9], diabetes long-term complication hospitalizations [WAS=70.3; ME=59.1], and diabetes mortality (underlying cause) per 100,000 population [WAS=33.9; ME=20.8]*</li> </ul>					
	<ul> <li>More children with unconfirmed elevated blood lead levels (% among those screened) [WAS=5.3%; ME=4.2%]*</li> </ul>					
	<ul> <li>High incidence of past or present hepatitis C virus (HCV) per 100,000 population [WAS=125.8; ME=107.1]</li> </ul>					
	<ul> <li>High pertussis incidence per 100,000 population [WAS=125.8; ME=41.9]</li> </ul>					

Health Issues - Surveillance Data					
Health Successes	Health Challenges				
	<ul> <li>Higher rates of death from unintentional and undetermined intent poisoning [WAS=15.6; ME=11.1], unintentional falls [WAS=9.8; ME=6.8], and motor vehicle traffic crashes per 100,000 population [WAS=19.5; ME=10.8]*</li> </ul>				
	<ul> <li>More deaths from firearms [WAS=17.0; ME=9.2]* and suicides [WAS=21.1; ME=15.2] per 100,000 population and higher violent crime rate per 100,000 population [WAS=174.0; ME=125.0]</li> </ul>				
<ul> <li>A higher percentage of high school students felt sad/hopeless for two weeks in a row [WAS=28.7 ME=24.3%]</li> </ul>					
<ul> <li>Higher rates of live births to 15-19 year olds per 1,000 population [WAS=28.7; ME=20.5]</li> </ul>					
<ul> <li>More substance and alcohol use and abuse compared to the state. This includes:</li> </ul>					
	<ul> <li>High alcohol-induced mortality per 100,000 population [WAS=11.0; ME=8.0]</li> </ul>				
	<ul> <li>High drug-induced mortality per 100,000 population [WAS=18.7; ME=12.4]</li> </ul>				
	<ul> <li>More chronic heavy drinking among adults than nationally [WAS=8.3%; U.S.=6.2%]</li> </ul>				
	<ul> <li>More binge drinking of alcoholic beverages among high school students) [WAS=17.8%; ME=14.8%]</li> </ul>				
	<ul> <li>More past-30-day inhalant use among high school students [WAS=4.8%; ME=3.2%]</li> </ul>				
	<ul> <li>High prescription Monitoring Program opioid prescriptions (days supply per population) [WAS=9.0; ME=6.8]</li> </ul>				
	<ul> <li>More drug-affected baby referrals received as a percentage of all live births [WAS=13.9%; ME=7.8%]</li> </ul>				

Asterisk (\*) indicates a statistically significant difference between Washington County and Maine All rates are per 100,000 population unless otherwise noted.

Table 23 summarizes the results of the health issues questions in the stakeholder survey for Washington County. It includes a summary of the biggest health challenges from the perspective of stakeholders who work in and represent communities in the county. The table also shares stakeholders' knowledge of the assets and resources available and those that are lacking but needed in the county to address the biggest health challenges.

Table 23.	Priority	Health	Issue	Challenges	and	Resources	for	Washington	County-
Stakehold	er Survey	Respons	ses						

Stakeholder Input - Stakeholder Survey Responses <sup>23</sup>				
Community Challenges	Community Resources			
Biggest health issues in Washington County according to stakeholders (% of	Assets Needed to Address Challenges:			
those rating issue as a major or critical problem in their area).	• <b>Drug and alcohol abuse:</b> Greater access to drug/alcohol treatments; greater access to substance abuse prevention programs; free or low-cost treatments for			
<ul> <li>Drug and alcohol abuse (91%)</li> <li>Obesity (82%)</li> </ul>	the uninsured; more substance abuse treatment providers; additional therapeutic programs			
<ul> <li>Tobacco use (77%)</li> <li>Mental health (76%)</li> <li>Cardiovascular diseases (76%)</li> </ul>	• <b>Obesity:</b> Greater access to affordable and healthy food; more programs that support low income families			
<ul> <li>Cardiovascular diseases (76%)</li> </ul>	• <b>Mental health:</b> More mental health professionals; more community-based services; better funding and support; greater access to inpatient care; readily available information about resources; transitional programs			
	Assets Available in County/State:			
	• Drug and alcohol abuse: Maine Alcoholics Anonymous; Substance Abuse Hotlines; Office of Substance Abuse and Mental Health Services			
	• <b>Obesity:</b> Public gyms; farmers markets; Maine SNAP-ED Program; school nutrition programs; public walking and biking trails; Healthy Maine Partnerships; Let's Go! 5-2- 1-0			
	<ul> <li>Mental health: Mental health/counseling providers and programs</li> </ul>			
	• Cardiovascular diseases: Hospitals; Primary Care Providers; YMCA's (Public gyms); Education programs			

<sup>&</sup>lt;sup>23</sup> Results are from the Maine Shared Community Health Needs Assessment Stakeholder Survey, conducted in May-June, 2015.

Table 24 presents a summary of the major health strengths and challenges that affect the health of Washington County residents. Data come from a comprehensive analysis of available surveillance data (see Table 28 for a full list of the health indicators and factors included in this project). Two criteria were used to select the factors presented in this table. **Statistically significant differences**, using a 95 percent confidence level, between the county and state are noted with an asterisk (\*) after the indicator. A **rate ratio** was also calculated to compare the relative difference between the county and state. Indicators where the county was 15 percent or more above or below the state average are included in this table.

Неа	alth Factors – Surveillance Data
Health Factor Strengths	Health Factor Challenges
<ul> <li>Fewer individuals who are unable to obtain or delay obtaining necessary medical care due to cost [WAS=9.6%; U.S.=15.3%]</li> <li>Higher percentage of lead</li> </ul>	<ul> <li>Washington County has a number of socioeconomic challenges when compared to the rest of the state, including: <ul> <li>A higher unemployment rate [WAS=8.4%; ME=5.7%]</li> <li>More individuals living in poverty [WAS=19.5%; ME=13.6%]* and children living in poverty [WAS=26.2%; ME=18.5%]*</li> <li>Lower median household income [WAS=\$37,236; ME=\$48,453]</li> </ul> </li> </ul>
screening among children age 12-23 months [WAS=79.5%; ME=49.2%]*	<ul> <li>Higher percentage of the population is uninsured [WAS=13.7%; ME=11.2%]* and fewer adults have a usual primary care provider [WAS=83.6%; ME=87.7%]*</li> </ul>
and children age 24-35 months [WAS=50.5%; ME=27.6%]*	<ul> <li>Fewer adults ages 18-34 with visits to a dentist in the past 12 months [WAS=56.9%; ME=65.3%]*</li> </ul>
	<ul> <li>Females age 50+ less likely to have a mammogram in the past 2 years [WAS=70.0%; ME=82.1%]*</li> </ul>
	<ul> <li>Lower percentage of adults [WAS=75.8%; ME=85.2%]* and high school students always wear seatbelts [WAS=42.6%; ME=61.6%]*</li> </ul>
	<ul> <li>More adults with a sedentary lifestyle – no leisure-time physical activity in past month [WAS=27.9%; ME=22.4%]*</li> </ul>
	<ul> <li>Among high school students, less fruit and vegetable consumption [WAS=14.0%; ME=16.8%], more soda/sports drink consumption [WAS=34.7%; ME=26.2%]* and higher levels of obesity [WAS=15.9%; ME=12.7%]</li> </ul>
	<ul> <li>Higher percentage of adults are current smokers [WAS=28.8%; ME=20.2%]</li> </ul>
	<ul> <li>More current smoking [WAS=19.7%; ME=12.9%] and current tobacco use among high school students [WAS=23.6%; ME=18.2%] and higher levels of secondhand smoke exposure among high school students [WAS=53.2%; ME=38.3%]*</li> </ul>

 
 Table 24. Priority Health Factor Strengths and Challenges for Washington County-Surveillance Data

Asterisk (\*) indicates a statistically significant difference between Washington County and Maine. All rates are per 100,000 population unless otherwise noted. Table 25 summarizes the results of the health factor questions in the stakeholder survey for Washington County. It includes a summary of the health factors that cause the biggest challenges from the perspective of stakeholders who work in and represent communities in the county. The table also shares stakeholders' knowledge of the assets and resources available and those that are lacking but needed in the county to address the biggest health challenges.

Table 25. Priority	<sup>r</sup> Health	Factor	Challenges	and	Resources	for	Washington	County-
Stakeholder Respo	nses							

Stakeholder Input- Stakeholder Survey Responses <sup>24</sup>					
Community Challenges	Community Resources				
Biggest health factors leading to poor health outcomes in Washington County according to stakeholders (% of those rating factor as a major or critical problem in their area).	<ul> <li>Assets Needed to Address Challenges:</li> <li>Poverty: Greater economic development; increased mentoring services; more skills trainings; more employment opportunities at livable wages; better transportation; better education</li> </ul>				
<ul> <li>Poverty (89%)</li> <li>Employment (82%)</li> <li>Health care insurance (78%)</li> <li>Access to behavioral care/mental health</li> </ul>	<ul> <li>Employment: More job creations; more training; more employment opportunities at livable wages; greater economic development; more funding for education</li> </ul>				
<ul> <li>Access to behavioral care/mental health care (76%)</li> <li>Health literacy (74%)</li> </ul>	• Health care insurance: Expansion of Medicaid; making insurance more affordable; universal health care; more stable health care system				
	• Access to behavioral care/mental health care: Better access to behavioral/mental health care for the uninsured; full behavioral/mental health integration at hospital and primary care levels; expand behavioral/mental health agencies to more rural areas; more hospital beds for mentally ill patients				
	Assets Available in County/State:				
	<ul> <li>Poverty: General Assistance; other federal, state and local programs</li> </ul>				
	• Employment: Adult education centers; career centers				
	<ul> <li>Health care insurance: MaineCare; ObamaCare (Affordable Care Act); Free care</li> </ul>				
	<ul> <li>Access to behavioral care/mental health care: Behavioral/mental health agencies</li> </ul>				
	<ul> <li>Health literacy: Hospital systems; primary care providers; social service agencies</li> </ul>				


<sup>&</sup>lt;sup>24</sup> Results are from the Maine Shared Community Health Needs Assessment Stakeholder Survey, conducted in May-June, 2015.

## County Health Rankings & Roadmaps

Each year, the University of Wisconsin Health Institute and Robert Wood Johnson Foundation produce *The County Health Rankings & Roadmaps* for every county in the U.S. The annual reports measure the social, economic, environmental and behavioral factors that influence health. These factors are quantified using indicators such as high school graduation rates, obesity, smoking, unemployment, access to healthy foods, the quality of air and water, income and teen births, to name a few. The rankings weight and score the sets of indicators to provide county comparisons within each state. For more information: www.countyhealthrankings.org

For this analysis, the 2015 rankings data for each of Maine's 16 counties is displayed in the graphic used by the University of Wisconsin to show how all of the factors ultimately affect community health. The comparison across counties provides insight into county health status. In Maine, the county ranked as "#1" on a particular health issue, is the healthiest in that measure, "#16" is the least healthy. The data for the underlying health measures are those used by the University of Wisconsin in its 2015 report and may not always match the data shown in other sections of this report due to timing or use of different indicators.

In interpreting the rankings for each county, it is important to keep in mind the underlying health measures. Because of the forced ranking, one county is always the "healthiest" and one is always the "least healthy." The comparisons are helpful in understanding differences, but it is important to look past the assignment of rank to understand the underlying issues and opportunities and their relative importance in the region.



#### Washington County

## Stakeholder Survey Findings

#### Table 26. Stakeholder Survey Results for Washington County and Maine

## Detailed Findings from SHNAPP Stakeholder Survey, June 2015

Survey Questions and Top Responses		
	Washington County	Maine
Demographics		
Which of the following sectors best describes your role or organization? (12 choices, picked 1)		
Number of Respondents	n=133	n=1639
Medical care provider	23%	22%
Other non-profit or social service agency	15%	14%
Other	17%	13%
Public health	8%	11%
Business owner or employee	8%	9%
Educator	11%	8%
Other type of health care organization	10%	8%
Behavioral/mental health provider	5%	6%
Local government	2%	4%
Other governmental agency	1%	3%
Youth-serving organization	2%	2%
Faith-based organization	1%	1%
Do you work for or represent: (5 choices, picked 1)		
None of the above	51%	49%
Hospital/Health-care system	40%	38%
Local public health agency	7%	10%
Maine CDC	2%	3%
Tribal health	0%	<1%
Please identify the type of geographical area that you primarily serve? (6	choices, picked 1)	
Town or region	31%	27%
Hospital/Health service area	28%	26%
Statewide	1%	22%
County	31%	18%
Other area	3%	4%
Public health district	7%	3%
Does your organization work with specific groups of people or population of, or experiencing, higher rates of health risk or poorer health outcomes within your area?	•	•
Yes	23%	24%
Somewhat	36%	47%
No	41%	29%

Detailed Findings from SHNAPP Stakeholder Surv	ey, June 2015	;
Survey Questions and Top Responses		
	Washington County	Maine
If "Yes" or "Somewhat" to Q4: To which of the following populations does	your organization	directly
provide resources to address their needs? (select all that apply)		
Number of Respondents	n=78	n=1159
Low-income, including those below the federal poverty limit, or defined as low-income by some other definition	76%	77%
Medically-underserved - including uninsured and under-insured	49%	63%
People with disabilities - physical, mental, or intellectual	42%	58%
Very rural and/or geographically isolated people	67%	47%
Less than a high school education and/ or low literacy (low reading or math skills)	36%	47%
Women	32%	44%
Limited or no English proficiency	24%	38%
Gay, lesbian, bisexual or transgendered people	27%	36%
Deaf and hard of hearing people	23%	35%
Military veterans	24%	34%
Refugees/immigrants	10%	28%
Racial/ethnic minority populations	19%	27%
Members of any federally recognized tribe	36%	25%
Specific age group	24%	21%
Other	14%	15%
Don't know	8%	5%
Overall, to what degree to you feel the health needs of your area are being		
Number of Respondents	n=133	n=1639
Not addressed at all	5%	<1%
Mostly unaddressed	17%	10%
Somewhat addressed	37%	55%
Mostly addressed	20%	30%
Completely addressed	8%	2%
Don't know	14%	2%
Health Issues and Factors		
Please rate the following health issues based on how you feel they impact residents in your area. ( <i>Percentage of stakeholders in county who rated iss</i>		
problem in their area)		
Number of Respondents	n=133	n=1639
Family Health		
Adolescent health	32%	25%
Child developmental issues	44%	34%
Childhood obesity	49%	58%
Elder health	64%	55%
Infant mortality	6%	4%

Detailed Findings from SHNAPP Stakeholder	Survey, June 2015	
Survey Questions and Top Respor	nses	
	Washington County	Maine
Maternal and child health	31%	23%
Chronic Diseases		
Cancer	74%	50%
Cardiovascular disease	76%	63%
Depression	74%	67%
Diabetes	74%	63%
Musculoskeletal diseases	40%	28%
Neurological diseases	38%	35%
Obesity	82%	78%
Respiratory disease	69%	60%
Infectious Diseases		
Infectious diseases	25%	22%
Sexually transmitted diseases/HIV/AIDS	14%	13%
Health Behaviors		
Drug and alcohol abuse	91%	80%
Physical activity and nutrition	74%	69%
Tobacco use	77%	63%
Other Health Issues	· · ·	
Lead poisoning and other environmental health issues	23%	17%
Mental health	76%	71%
Oral health	67%	53%
Suicide and self-harm	46%	37%
Unintentional injury	48%	34%
Violence	42%	38%
"Don't know" responses not includ	ed	
Please indicate how much of a problem each of the following health poor health outcomes for residents. ( <i>Percentage of stakeholders in or critical problem in their area</i> )	-	
Number of Respondents	n=133	n=1639
Economic Stability		
Employment	82%	64%
Food security	66%	58%
Housing stability	59%	57%
Poverty	89%	78%
Education		
Enrollment in higher education	41%	35%
Early childhood education/development	57%	43%
High school graduation	35%	31%
Language and literacy	39%	34%

Detailed Findings from SHNAPP Stakeholder Surv	ey, June 2015	
Survey Questions and Top Responses		
	Washington County	Maine
Social and Community Context		
Adverse childhood experiences	64%	56%
Civic participation	34%	30%
Incarceration or institutionalization	35%	35%
Social attitudes such as discrimination	46%	38%
Social support and interactions	53%	50%
Caregiver support	48%	46%
Health and Health Care		
Access to behavioral care/mental health care	76%	67%
Access to primary care	56%	39%
Access to other health care	57%	41%
Access to oral health	66%	56%
Health care insurance	78%	64%
Health literacy	74%	62%
Neighborhood and Built Environment	<u> </u>	
Access to healthy foods	58%	53%
Access to physical activity opportunities	59%	42%
Crime and violence	37%	27%
Environmental conditions	16%	12%
Quality of housing	38%	34%
Transportation	70%	67%
* "Don't know" responses not included		
Please rank each health issue according to how you think resources in your (1=highest priority and 8=lowest priority) (mean)	r area should be al	located.
Number of Respondents	n=92	n=1168
Risk factors that lead to poor health	2.98	3.08
Mental health - conditions that impact how people think, feel and act as they cope with life	3.54	3.49
Substance abuse behaviors, including excessive drinking, smoking, and other drug use	3.05	3.71
Community capacity - ability to sustain a high quality of life, including access to employment, education and housing	3.38	3.93
Chronic diseases, such as heart disease, cancer, diabetes, and asthma	3.80	4.05
Family health, including teen pregnancy, prenatal care, and healthy behaviors during pregnancy	4.32	4.81
Environmental issues - access to healthy foods, access to recreation, clean air, water, lead exposure, etc.	5.63	5.36
Injuries, intentional and unintentional	6.58	6.52

## Health Indicators Results from Secondary Data Sources

The county level summary of health indicators analyzed from secondary data sources is presented in the table below. Results are displayed for the county, state and U.S. (where available). County trends are presented in the column after the county data when available. Results are organized by health issue or category. Please note that age-adjusted rates are presented for all applicable indicators, with the exception of ambulatory care-sensitive conditions and infectious and sexually transmitted diseases (which are presented as crude rates). A detailed list of all data sources, years and notes for all indicators is presented in Table 28.

Indicates county is significantly better than state average (using a 95% confidence level).
 Indicates county is significantly worse than state average (using a 95% confidence level).
 + Indicates an improvement in the indicator over time at the county level (using a 95%)

confidence level)

- Indicates a worsening in the indicator over time at the county level (using a 95% confidence level)

*† Results may be statistically unreliable due to small numerator, use caution when interpreting. NA = Data not available.* 

Maine Shared CHNA Health Indicators	Year	Washington	Trend	Maine	U.S.
Demographics					
Total Population	2013	32,190		1,328,302	319 Mil
Population – % ages 0-17	2013	18.9%		19.7%	23.3%
Population – % ages 18-64	2013	59.7%		62.6%	62.6%
Population – % ages 65+	2013	21.4%		17.7%	14.1%
Population – % White	2013	92.0%		95.2%	77.7%
Population – % Black or African American	2013	0.6%		1.4%	13.2%
Population – % American Indian and Alaska Native	2013	5.1%		0.7%	1.2%
Population – % Asian	2013	0.5%		1.1%	5.3%
Population – % Hispanic	2013	1.7%		1.4%	17.1%
Population – % with a disability	2013	20.5%		15.9%	12.1%
Population density (per square mile)	2013	12.8		43.1	87.4
Socioeconomic Status Measures					
Adults and children living in poverty	2009-2013	19.5%	NA	13.6%	15.4%
Children living in poverty	2009-2013	26.2%	NA	18.5%	21.6%
High school graduation rate	2013-2014	89.4%	NA	86.5%	81.0%
Median household income	2009-2013	\$37,236	NA	\$48,453	\$53 <i>,</i> 046
Percentage of people living in rural areas	2013	100.0%	NA	66.4%	NA
Single-parent families	2009-2013	37.0%	NA	34.0%	33.2%
Unemployment rate	2014	8.4%	NA	5.7%	6.2%
65+ living alone	2009-2013	44.3%	NA	41.2%	37.7%
General Health Status					
Adults who rate their health fair to poor	2011-2013	20.4%		15.6%	16.7%

Table 27. Quantitative Health Indicators for Washington County, Maine and the U.S.

Indicates county is significantly better than state average (using a 95% confidence level).

Indicates county is significantly worse than state average (using a 95% confidence level).

Maine Shared Community Health Needs Assessment, 2016 page 41 www.maine.gov/SHNAPP/county-reports.shtml

Maine Shared CHNA Health Indicators	Year	Washington	Trend	Maine	U.S.
Adults with 14+ days lost due to poor mental health	2011-2013	12.6%	menta	12.4%	NA
Adults with 14+ days lost due to poor incitial health	2011-2013	14.4%		13.1%	NA
Adults with three or more chronic conditions	2011, 2013	31.6%		27.6%	NA
Mortality	2011, 2013	51.070		27.070	1177
Life expectancy (Female)	2012	80.1	NA	81.5	81.2
Life expectancy (Male)	2012	74.7	NA	76.7	76.4
Overall mortality rate per 100,000 population	2009-2013	875.0	NA	745.8	731.9
Access	2003 2013	075.0	nA.	745.0	751.5
Adults with a usual primary care provider	2011-2013	83.6%		87.7%	76.6%
Individuals who are unable to obtain or delay		00.070		0/11/0	/ 0.0/0
obtaining necessary medical care due to cost	2011-2013	9.6%		11.0%	15.3%
MaineCare enrollment	2015	40.3%	NA	27.0%	23.0%
Percent of children ages 0-19 enrolled in MaineCare	2015	61.5%	NA	41.8%	48.0%
Percent uninsured	2009-2013	13.7%	NA	10.4%	11.7%
Health Care Quality					
Ambulatory care-sensitive condition hospital					
admission rate per 100,000 population	2011	1,809.0	+	1,499.3	1457.5
Ambulatory care-sensitive condition emergency					
department rate per 100,000 population	2011	6,645.8	NA	4,258.8	NA
Oral Health					
Adults with visits to a dentist in the past 12 months	2012	56.9%	NA	65.3%	67.2%
MaineCare members under 18 with a visit to the		60 <b>F</b> 0(			
dentist in the past year	2014	60.5%	NA	55.1%	NA
Respiratory					
Asthma emergency department visits per 10,000	2009-2011	118.0		67.3	NIA
population	2009-2011	118.0		07.3	NA
COPD diagnosed	2011-2013	8.6%		7.6%	6.5%
COPD hospitalizations per 100,000 population	2011	187.7		216.3	NA
Current asthma (Adults)	2011-2013	10.2%		11.7%	9.0%
Current asthma (Youth 0-17)	2011-2013	10.9%†	NA	9.1%	NA
Pneumonia emergency department rate per 100,000	2011	1,429.8		719.9	NA
population	2011	1,429.0		/19.9	
Pneumonia hospitalizations per 100,000 population	2011	714.0	-	329.4	NA
Cancer					
Mortality – all cancers per 100,000 population	2007-2011	210.5	NA	185.5	168.7
Incidence – all cancers per 100,000 population	2007-2011	533.3	NA	500.1	453.4
Bladder cancer incidence per 100,000 population	2007-2011	24.0	NA	28.3	20.2
Female breast cancer mortality per 100,000	2007-2011	23.8	NA	20.0	21.5
population	2007-2011	23.0		20.0	21.3
Breast cancer late-stage incidence (females only) per	2007-2011	47.1	NA	41.6	43.7
100,000 population		.,			
Female breast cancer incidence per 100,000	2007-2011	122.7	NA	126.3	124.1
population					
Mammograms females age 50+ in past two years	2012	70.0%	NA	82.1%	77.0%
Colorectal cancer mortality per 100,000 population	2007-2011	14.9	NA	16.1	15.1
Colorectal late-stage incidence per 100,000	2007-2011	26.4	NA	22.7	22.9

Indicates county is significantly worse than state average (using a 95% confidence level).

Maine Shared Community Health Needs Assessment, 2016 page 42 www.maine.gov/SHNAPP/county-reports.shtml

Maine Shared CHNA Health Indicators	Year	Washington	Trend	Maine	U.S.
population	Tear	washington	menta	wianie	0.5.
Colorectal cancer incidence per 100,000 population	2007-2011	52.8	NA	43.5	42.0
Colorectal screening	2007 2011	66.3%	NA	72.2%	42.0 NA
Lung cancer mortality per 100,000 population	2007-2011	61.8	NA	54.3	46.0
Lung cancer incidence per 100,000 population	2007-2011	92.8	NA	75.5	58.6
Melanoma incidence per 100,000 population	2007-2011	11.1	NA	22.2	21.3
Pap smears females ages 21-65 in past three years	2007-2011	82.6%	NA	88.0%	78.0%
Prostate cancer mortality per 100,000 population	2012	29.1	NA	22.1	20.8
		143.8			
Prostate cancer incidence per 100,000 population	2007-2011	143.8	NA	133.8	140.8
Tobacco-related neoplasms, mortality per 100,000 population	2007-2011	46.3	NA	37.4	34.3
Tobacco-related neoplasms, incidence per 100,000 population	2007-2011	97.9	NA	91.9	81.7
Cardiovascular Disease	1	I			I
Acute myocardial infarction hospitalizations per 10,000 population	2010-2012	39.4		23.5	NA
Acute myocardial infarction mortality per 100,000 population	2009-2013	66.8	NA	32.2	32.4
Cholesterol checked every five years	2011. 2013	75.7%		81.0%	76.4%
Coronary heart disease mortality per 100,000 population	2009-2013	130.8	NA	89.8	102.6
Heart failure hospitalizations per 10,000 population	2010-2012	27.8		21.9	NA
Hypertension prevalence	2011, 2013	39.7%		32.8%	31.4%
High cholesterol	2011, 2013	45.9%		40.3%	38.4%
Hypertension hospitalizations per 100,000 population	2011	27.3		28.0	NA
Stroke hospitalizations per 10,000 population	2010-2012	22.5		20.8	NA
Stroke mortality per 100,000 population	2009-2013	27.4	NA	35.0	36.2
Diabetes	1	I			
Diabetes prevalence (ever been told)	2011-2013	10.4%		9.6%	9.7%
Pre-diabetes prevalence	2011-2013	3.4%†	NA	6.9%	NA
Adults with diabetes who have eye exam annually	2011-2013	NA	NA	71.2%	NA
Adults with diabetes who have foot exam annually	2011-2013	NA	NA	83.3%	NA
Adults with diabetes who have had an A1C test twice per year	2011-2013	NA	NA	73.2%	NA
Adults with diabetes who have received formal diabetes education	2011-2013	NA	NA	60.0%	55.8%
Diabetes emergency department visits (principal diagnosis) per 100,000 population	2011	294.1		235.9	NA
Diabetes hospitalizations (principal diagnosis) per 10,000 population	2010-2012	10.9		11.7	NA
Diabetes long-term complication hospitalizations	2011	70.3		59.1	NA
Diabetes mortality (underlying cause) per 100,000 population	2009-2013	33.9	NA	20.8	21.2
Environmental Health					
Children with confirmed elevated blood lead levels (% among those screened)	2009-2013	1.2%	NA	2.5%	NA

Indicates county is significantly worse than state average (using a 95% confidence level).

Maine Shared Community Health Needs Assessment, 2016 page 43 www.maine.gov/SHNAPP/county-reports.shtml

Maine Shared CHNA Health Indicators	Year	Washington	Trend	Maine	U.S.
Children with unconfirmed elevated blood lead levels			пспа	wante	0.5.
(% among those screened)	2009-2013	5.3%	NA	4.2%	NA
Homes with private wells tested for arsenic	2009, 2012	34.1%	NA	43.3%	NA
Lead screening among children age 12-23 months	2009-2013	79.5%	NA	49.2%	NA
Lead screening among children age 24-35 months	2009-2013	50.5%	NA	27.6%	NA
Immunization					
Adults immunized annually for influenza	2011-2013	42.0%		41.5%	NA
Adults immunized for pneumococcal pneumonia (ages 65 and older)	2011-2013	68.0%		72.4%	69.5%
Immunization exemptions among kindergarteners for philosophical reasons	2015	0.4%	NA	3.7%	NA
Two-year-olds up to date with "Series of Seven Immunizations" 4-3-1-3-3-1-4	2015	NA	NA	75.0%	NA
Infectious Disease					1
Hepatitis A (acute) incidence per 100,000 population	2014	0.0+	NA	0.6	0.4
Hepatitis B (acute) incidence per 100,000 population	2014	3.1†	NA	0.9	0.9
Hepatitis C (acute) incidence per 100,000 population	2014	6.3†	NA	2.3	0.7
Incidence of past or present hepatitis C virus (HCV) per 100,000 population	2014	125.8	NA	107.1	NA
Incidence of newly reported chronic hepatitis B virus (HBV) per 100,000 population	2014	0.0†	NA	8.1	NA
Lyme disease incidence per 100,000 population	2014	44.0†	NA	105.3	10.5
Pertussis incidence per 100,000 population	2014	125.8	NA	41.9	10.3
Tuberculosis incidence per 100,000 population	2014	0.0†	NA	1.1	3.0
STD/HIV					
AIDS incidence per 100,000 population	2014	0.0†	NA	2.1	8.4
Chlamydia incidence per 100,000 population	2014	157.2	NA	265.5	452.2
Gonorrhea incidence per 100,000 population	2014	15.7†	NA	17.8	109.8
HIV incidence per 100,000 population	2014	0.0+	NA	4.4	11.2
HIV/AIDS hospitalization rate per 100,000 population	2011	23.3		21.4	NA
Syphilis incidence per 100,000 population	2014	6.3†	NA	1.6	19.9
Intentional Injury					
Domestic assaults reports to police per 100,000 population	2013	394.6	NA	413.0	NA
Firearm deaths per 100,000 population	2009-2013	17.0	NA	9.2	10.4
Intentional self-injury (Youth)	2013	NA	NA	17.9%	NA
Lifetime rape/non-consensual sex (among females)	2013	NA	NA	11.3%	NA
Nonfatal child maltreatment per 1,000 population	2013	NA	NA	14.6	9.1
Reported rape per 100,000 population	2013	9.3†	NA	27.0	25.2
Suicide deaths per 100,000 population	2009-2013	21.1	NA	15.2	12.6
Violence by current or former intimate partners in past 12 months (among females)	2013	NA	NA	0.8%	NA
Violent crime rate per 100,000 population	2013	174.0	NA	125.0	368
Unintentional Injury					
Always wear seatbelt (Adults)	2013	75.8%		85.2%	NA
Always wear seatbelt (High School Students)	2013	42.6%		61.6%	54.7%

Indicates county is significantly worse than state average (using a 95% confidence level).

Maine Shared Community Health Needs Assessment, 2016 page 44 www.maine.gov/SHNAPP/county-reports.shtml

Maine Shared CHNA Health Indicators	Year	Washington	Trend	Maine	U.S.
Traumatic brain injury related emergency department visits (all intents) per 10,000 population	2011	67.9	NA	81.4	NA
Unintentional and undetermined intent poisoning deaths per 100,000 population	2009-2013	15.6	NA	11.1	13.2
Unintentional fall related deaths per 100,000 population	2009-2013	9.8	NA	6.8	8.5
Unintentional fall related injury emergency department visits per 10,000 population	2011	415.6	NA	361.3	NA
Unintentional motor vehicle traffic crash related deaths per 100,000 population	2009-2013	19.5	NA	10.8	10.5
Occupational Health					
Deaths from work-related injuries (number)	2013	NA	NA	19.0	4,585
Nonfatal occupational injuries (number)	2013	249.0	NA	13,205.0	NA
Mental Health					
Adults who have ever had anxiety	2011-2013	20.1%		19.4%	NA
Adults who have ever had depression	2011-2013	23.0%		23.5%	18.7%
Adults with current symptoms of depression	2011-2013	11.4%		10.0%	NA
Adults currently receiving outpatient mental health treatment	2011-2013	20.1%		17.7%	NA
Co-morbidity for persons with mental illness	2011, 2013	40.0%	NA	35.2%	NA
Mental health emergency department rates per 100,000 population	2011	2,061.2		1,972.1	NA
Sad/hopeless for two weeks in a row (High School Students)	2013	28.7%		24.3%	29.9%
Seriously considered suicide (High School Students)	2013	15.4%		14.6%	17.0%
Physical Activity, Nutrition and Weight			1		
Fewer than two hours combined screen time (High					
School Students)	2013	NA	NA	33.9%	NA
Fruit and vegetable consumption (High School Students)	2013	14.0%	NA	16.8%	NA
Fruit consumption among Adults 18+ (less than one serving per day)	2013	35.4%	NA	34.0%	39.2%
Met physical activity recommendations (Adults)	2013	45.9%		53.4%	50.8%
Physical activity for at least 60 minutes per day on five of the past seven days (High School Students)	2013	41.6%	NA	43.7%	47.3%
Sedentary lifestyle – no leisure-time physical activity in past month (Adults)	2011-2013	27.9%		22.4%	25.3%
Soda/sports drink consumption (High School Students)	2013	34.7%	NA	26.2%	27.0%
Vegetable consumption among Adults 18+ (less than one serving per day)	2013	19.7%	NA	17.9%	22.9%
Obesity (Adults)	2013	30.4%		28.9%	29.4%
Obesity (High School Students)	2013	15.9%		12.7%	13.7%
Overweight (Adults)	2013	35.0%		36.0%	35.4%
Overweight (High School Students)	2013	17.1%		16.0%	16.6%
Pregnancy and Birth Outcomes					
Children with special health care needs	2009-2010	NA	NA	23.6%	19.8%

Indicates county is significantly worse than state average (using a 95% confidence level).

Maine Shared Community Health Needs Assessment, 2016 page 45 www.maine.gov/SHNAPP/county-reports.shtml

Maine Shared CHNA Health Indicators	Year	Washington	Trend	Maine	U.S.
Infant deaths per 1,000 live births	2003-2012	4.3	NA	6.0	6.0
Live births for which the mother received early and adequate prenatal care	2010-2012	84.3%	NA	86.4%	84.8%
Live births to 15-19 year olds per 1,000 population	2010-2012	28.7	NA	20.5	26.5
Low birth weight (<2500 grams)	2010-2012	4.7%	NA	6.6%	8.0%
Substance and Alcohol Abuse					
Alcohol-induced mortality per 100,000 population	2009-2013	11.0	NA	8.0	8.2
Binge drinking of alcoholic beverages (High School Students)	2013	17.8%		14.8%	20.8%
Binge drinking of alcoholic beverages (Adults)	2011-2013	17.0%		17.4%	16.8%
Chronic heavy drinking (Adults)	2011-2013	8.3%		7.3%	6.2%
Drug-affected baby referrals received as a percentage of all live births	2014	13.9%	NA	7.8%	NA
Drug-induced mortality per 100,000 population	2009-2013	18.7	NA	12.4	14.6
Emergency medical service overdose response per 100,000 population	2014	389.8	NA	391.5	NA
Opiate poisoning (ED visits) per 100,000 population	2009-2011	20.8		25.1	NA
Opiate poisoning (hospitalizations) per 100,000 population	2009-2011	11.1		13.2	NA
Past-30-day alcohol use (High School Students)	2013	28.2%		26.0%	34.9%
Past-30-day inhalant use (High School Students)	2013	4.8%		3.2%	NA
Past-30-day marijuana use (Adults)	2011-2013	8.8%†	NA	8.2%	NA
Past-30-day marijuana use (High School Students)	2013	23.6%		21.6%	23.4%
Past-30-day nonmedical use of prescription drugs (Adult)	2011-2013	0.3%†	NA	1.1%	NA
Past-30-day nonmedical use of prescription drugs (High School Students)	2013	4.2%	+	5.6%	NA
Prescription Monitoring Program opioid prescriptions (days supply/pop)	2014-2015	9.0	NA	6.8	NA
Substance-abuse hospital admissions per 100,000 population	2011	360.0	-	328.1	NA
Tobacco Use					
Current smoking (Adults)	2011-2013	28.8%		20.2%	19.0%
Current smoking (High School Students)	2013	19.7%		12.9%	15.7%
Current tobacco use (High School Students)	2013	23.6%	NA	18.2%	22.4%
Secondhand smoke exposure (Youth)	2013	53.2%		38.3%	NA



Indicates county is significantly better than state average (using a 95% confidence level). Indicates county is significantly worse than state average (using a 95% confidence level).

		2015	
Indicator	Data Source	Year(s)	Other Notes
Demographics			
Population	U.S. Census	2013	2013 data was used for all age, racial and ethnic groups.
Population with a disability	U.S. Census	2011-2013	Adults reporting any one of the six disability types are considered to have a disability: hearing difficulty, vision difficulty, cognitive difficulty, ambulatory difficulty, self-care difficulty, independent living difficulty.
Population density	U.S. Census	2010	Based on 2010 U.S. Census population.
Socioeconomic Status Measures			
Adults and children living in poverty	U.S. Census	2009-2013	The poverty status of the household is determined by the poverty status of the householder. Households are classified as poor when the total income of the householder's family is below the appropriate poverty threshold. The American Community Survey measures poverty in the previous 12 months instead of the previous calendar year.
Children living in poverty	U.S. Census	2009-2013	The poverty status of the household is determined by the poverty status of the householder. Households are classified as poor when the total income of the householder's family is below the appropriate poverty threshold. The American Community Survey measures poverty in the previous 12 months instead of the previous calendar year.
High school graduation rate	Maine Dept. of Education	2013-14 School Year	Proportion of students who graduate with a regular diploma four years after starting ninth grade. Graduation rates include all public schools and all private schools that have 60% or more publicly funded students.
Median household income	U.S. Census	2009-2013	In 2013 inflation-adjusted dollars. This includes the income of the householder and all other individuals 15 years old and older in the household, whether they are related to the householder or not.
Percentage of people living in rural areas	U.S. Census	2012	The urban/rural categories used in this analysis were defined by the New England Rural Health Roundtable available in Rural Data For Action 2nd Edition: http://www.newenglandruralhealth.org/rural_data
Single-parent families	U.S. Census	2009-2013	Families consist of a householder and one or more other people related to the householder by birth, marriage, or adoption. They do not include same-sex married couples even if the marriage was performed in a state issuing marriage certificates for same-sex couples. "Householder without a spouse present" is defined as a male householder without a wife present or a female householder without a husband present.

# Table 28. List of Data Sources and Years for Quantitative Health Indicators Maine Shared Community Health Needs Assessment Data Sources

Maine Shared Community Health Needs Assessment Data Sources					
		2015			
Indicator	Data Source	Year(s)	Other Notes		
Unemployment rate	Bureau of Labor Statistics	2014	Unemployment rate of the civilian noninstitutionalized population averaged for the full year of 2014.		
65+ living alone	U.S. Census	2009-2013	Estimated number of one-person households with a person 65 years and older.		
General Health Status					
Adults who rate their health fair to poor	BRFSS	2011-2013	Adults rating their health as fair or poor vs. excellent, very good or good.		
Adults with 14+ days lost due to poor mental health	BRFSS	2011-2013	Now thinking about your mental health, which includes stress, depression and problems with emotions, for how many days during the past 30 days was your mental health not good?		
Adults with 14+ days lost due to poor physical health	BRFSS	2011-2013	Now thinking about your physical health, which includes physical illness and injury, for how many days during the past 30 days was your physical health not good?		
Adults with three or more chronic conditions	BRFSS	2011, 2013	Chronic conditions available in 2013 BRFSS: arthritis, asthma, cancer, cardiovascular disease, chronic kidney disease, chronic obstructive pulmonary disease (COPD), coronary heart disease, diabetes, hypertension, high cholesterol, obesity.		
Mortality					
Life expectancy (Female)	National Center for Health Statistics	2012	Life expectancy at birth.		
Life expectancy (Male)	National Center for Health Statistics	2012	Life expectancy at birth.		
Overall mortality rate per 100,000 population	DRVS	2009-2013	All deaths are defined as deaths in which the underlyi ng cause of death was coded as ICD-10 any listed.		
Access					
Adults with a usual primary care provider	BRFSS	2011-2013	Adults that have one or more person they think of as their personal doctor or health care provider.		
Individuals who are unable to obtain or delay obtaining necessary medical care due to cost	BRFSS	2011-2013	Adults reporting that there was a time during the last 12 months when they needed to see a doctor but could not because of the cost.		
MaineCare enrollment	MaineCare	2015	The number and percent of individuals participating in MaineCare. These data are reported as of April 2015. Percentages calculated based on the 2014 US Census population estimates. Individuals are reported by county of residence at the end of the SFY or the end of participation in the program. Figures exclude individuals who were nonresidents or who were out of state.		
Percent of children ages 0-19 enrolled in MaineCare	MaineCare	2015	The number and percent of individuals participating in MaineCare. These data are reported as of April 2015. Individuals are reported by county of residence at the end of the SFY or the end of participation in		

Maine Shared Community Health Needs Assessment Data Sources					
		2015			
Indicator	Data Source	Year(s)	Other Notes		
			the program. Figures exclude individuals who were		
			nonresidents or who were out of state.		
Percent uninsured	U.S. Census	2009-2013	Estimated number of Maine people who do not currently have health insurance.		
Health Care Quality					
Ambulatory care-sensitive condition hospital admission rate per 100,000 population	MHDO	2011	PQI = Prevention Quality Indicators, a set of measures that can be used with hospital inpatient discharge data to identify quality of care for ambulatory care-sensitive conditions. Additional information at: AHRQ Quality Indicators, Version 4.4,		
			Agency for Healthcare Research and Quality: U.S. Department of Health and Human Services. http://www.qualityindicators.ahrq.gov.		
Ambulatory care-sensitive condition emergency department rate per 100,000 population	MHDO	2011	PQI = Prevention Quality Indicators, a set of measures that can be used with hospital inpatient discharge data to identify quality of care for ambulatory care-sensitive conditions. Additional information at: AHRQ Quality Indicators, Version 4.4, Agency for Healthcare Research and Quality: U.S. Department of Health and Human Services. http://www.qualityindicators.ahrq.gov.		
Oral Health					
Adults with visits to a dentist in the past 12 months	BRFSS	2012	Adults who last visited the dentist or a dental clinic for any reason in the past 12 months.		
MaineCare members under 18 with a visit to the dentist in the past year	Maine Care	2014	Total members younger than 18 with dental claims during calendar year 2014 was 67,871. Of those, only 61,948 had eligibility as of April 2015. Members were younger than 18 on date of service, but some turned 18 by April 2015.		
Respiratory					
Asthma emergency department visits per 10,000 population	MHDO	2009-2011	ICD-9 CM - 493		
COPD diagnosed	BRFSS	2011-2013	Adults that have been told by a doctor, nurse or health professional that they have COPD chronic obstructive pulmonary disease, emphysema, or chronic bronchitis.		
COPD hospitalizations per 100,000 population	MHDO	2011	ICD-9 CM - 490, 491, 492, 494, 496		
Current asthma (Adults)	BRFSS	2011-2013	Adults that have been told by a doctor, nurse or health professional that they had asthma and that they still have asthma.		
Current asthma (Youth 0-17)	BRFSS	2011-2013	Children that have been told by a doctor, nurse or health professional that they had asthma and that they still have asthma.		
Pneumonia emergency department rate per 100,000 population	MHDO	2011	ICD-9 CM - 480-486		

Maine Shared Community Health Needs Assessment Data Sources					
		2015			
Indicator	Data Source	Year(s)	Other Notes		
Pneumonia hospitalizations per 100,000 population	MHDO	2011	ICD-9 CM - 480-486		
Cancer					
Mortality – all cancers per 100,000 population	MCR	2007-2011	All cancer: SEER Cause of Death Recode: 20010- 37000 (which include ICD-10 codes: C00-C97).		
Incidence – all cancers per 100,000 population	MCR	2007-2011	All cancer: SEER Site Recode: 20010-37000 (which include ICD-O-3 codes: C00-C797).		
Bladder cancer incidence per 100,000 population	MCR	2007-2011	Cancer Incidence: The number of people who develop cancer (new cancer cases) during a specified period of time in a specified population. Incidence case definitions exclude histologies consistent with Kaposi sarcoma and mesothelioma, where applicable.		
Female breast cancer mortality per 100,000 population	MCR	2007-2011	Cancer Deaths: Deaths with malignant cancer as the underlying cause of death.		
Breast cancer late-stage incidence (females only) per 100,000 population	Maine Cancer Registry (MCR)	2007-2011	Cancer Incidence: The number of people who develop cancer (new cancer cases) during a specified period of time in a specified population. Incidence case definitions exclude histologies consistent with Kaposi sarcoma and mesothelioma, where applicable.		
Female breast cancer incidence per 100,000 population	MCR	2007-2011	Cancer Incidence: The number of people who develop cancer (new cancer cases) during a specified period of time in a specified population. Incidence case definitions exclude histologies consistent with Kaposi sarcoma and mesothelioma, where applicable.		
Mammograms females age 50+ in past two years	BRFSS	2012	Females ages 50 years and older who reported they had a mammogram within the past 2 years.		
Colorectal cancer mortality per 100,000 population	MCR	2007-2011	Cancer Deaths: Deaths with malignant cancer as the underlying cause of death.		
Colorectal late-stage incidence per 100,000 population	MCR	2007-2011	Cancer Incidence: The number of people who develop cancer (new cancer cases) during a specified period of time in a specified population. Incidence case definitions exclude histologies consistent with Kaposi sarcoma and mesothelioma, where applicable.		
Colorectal cancer incidence per 100,000 population	MCR	2007-2011	Cancer Incidence: The number of people who develop cancer (new cancer cases) during a specified period of time in a specified population. Incidence case definitions exclude histologies consistent with Kaposi sarcoma and mesothelioma, where applicable.		
Colorectal screening	BRFSS	2012	Adults ages 50 years and older who reported that they had a home blood stool test (e.g., FOBT or FIT) within the past year OR sigmoidoscopy within the past 5 years and home blood stool test within the past 3 years OR a colonoscopy within the past 10 years.		

# Maine Shared Community Health Needs Assessment Data Sources

		2015	
Indicator	Data Source	Year(s)	Other Notes
Lung cancer mortality per 100,000 population	MCR	2007-2011	Cancer Deaths: Deaths with malignant cancer as the underlying cause of death.
Lung cancer incidence per 100,000 population	MCR	2007-2011	Cancer Incidence: The number of people who develop cancer (new cancer cases) during a specified period of time in a specified population. Incidence case definitions exclude histologies consistent with Kaposi sarcoma and mesothelioma, where applicable.
Melanoma incidence per 100,000 population	MCR	2007-2011	Cancer Incidence: The number of people who develop cancer (new cancer cases) during a specified period of time in a specified population. Incidence case definitions exclude histologies consistent with Kaposi sarcoma and mesothelioma, where applicable.
Pap smears females ages 21-65 in past three years	BRFSS	2012	Females with intact cervix, that have received a pap smear within the past three years.
Prostate cancer mortality per 100,000 population	MCR	2007-2011	Cancer Deaths: Deaths with malignant cancer as the underlying cause of death.
Prostate cancer incidence per 100,000 population	MCR	2007-2011	Cancer Incidence: The number of people who develop cancer (new cancer cases) during a specified period of time in a specified population. Incidence case definitions exclude histologies consistent with Kaposi sarcoma and mesothelioma, where applicable.
Tobacco-related neoplasms, mortality per 100,000 population	MCR	2007-2011	Cancer Deaths: Deaths with malignant cancer as the underlying cause of death.
Tobacco-related neoplasms, incidence per 100,000 population	MCR	2007-2011	Cancer Incidence: The number of people who develop cancer (new cancer cases) during a specified period of time in a specified population. Incidence case definitions exclude histologies consistent with Kaposi sarcoma and mesothelioma, where applicable.
Cardiovascular Disease			
Acute myocardial infarction hospitalizations per 10,000 population	MHDO	2010-2012	ICD-9 CM - 410
Acute myocardial infarction mortality per 100,000 population	Maine CDC Vital Records	2009-2013	ICD-10 I21-I22
Cholesterol checked every five years	BRFSS	2011. 2013	Adults reporting that they last had their blood choices choices the choice choice choice choices choices and the choice c
Coronary heart disease mortality per 100,000 population	Maine CDC Vital Records	2009-2013	ICD-10 I20-I25
Heart failure hospitalizations per 10,000 population	MHDO	2010-2012	ICD-9 CM - 428
Hypertension prevalence	BRFSS	2011, 2013	Adults who have ever been told by a doctor, nurse, or other health professional that they have high blood pressure.

Maine Shared Community Health Needs Assessment Data Sources						
		2015				
Indicator	Data Source	Year(s)	Other Notes			
High cholesterol	BRFSS	2011, 2013	Adults who have been told by a doctor or other health professional that their blood cholesterol is high.			
Hypertension hospitalizations per 100,000 population	MHDO	2011	ICD-9 CM - 401, 402, 403, 404			
Stroke hospitalizations per 10,000 population	MHDO	2010-2012	ICD-9 CM - 430-438			
Stroke mortality per 100,000 population	Maine CDC Vital Records	2009-2013	ICD-10 I60-I69			
Diabetes						
Diabetes prevalence (ever been told)	BRFSS	2011-2013	Adults that have ever been told by a doctor or other health professional that they have diabetes.			
Pre-diabetes prevalence	BRFSS	2011-2013	Adults that have ever been told by a doctor or other health professional that they have pre-diabetes or borderline diabetes.			
Adults with diabetes who have eye exam annually	BRFSS	2011-2013	Adults with diabetes who report having an eye exam in which the pupils were dilated within the past year.			
Adults with diabetes who have foot exam annually	BRFSS	2011-2013	Adults with diabetes who report having a health professional check their feet for any sores or irritations within the past year.			
Adults with diabetes who have had an A1C test twice per year	BRFSS	2011-2013	Adults who have had a doctor, nurse, or other health professional checked them for "A one C" in the past 12 months.			
Adults with diabetes who have received formal diabetes education	BRFSS	2011-2013	Adults with diabetes who have ever taken a course or class in how to manage your diabetes themselves.			
Diabetes emergency department visits (principal diagnosis) per 100,000 population	MHDO	2011	ICD-9 CM - 250			
Diabetes hospitalizations (principal diagnosis) per 10,000 population	MHDO	2010-2012	ICD-9 CM - 250			
Diabetes long-term complication hospitalizations	MHDO	2011	Diabetes long-term complication hospitalizations are defined as hospitalizations of Maine residents for which diabetes long-term complication was the primary diagnosis, coded as ICD 9 - 25040, 25070, 25041, 25071, 25042, 25072, 25043, 25073, 25050, 25051, 25052, 25053, 25080, 25081, 25082, 25083, 25060, 25061, 25062, 25063, 25090, 25091, 25092.			
Diabetes mortality (underlying cause) per 100,000 population	Maine CDC Vital Records	2009-2013	ICD-10 E10-E14			
Environmental Health						
Children with confirmed elevated blood lead levels (% among those screened)	Maine CDC Lead Program	2009-2013	In 2012, CDC defined a reference level of 5 micrograms per deciliter ( $\mu$ g/dL) to identify children with elevated blood lead levels. These children are exposed to more lead than most children. For more information, visit: www.cdc.gov/nceh/lead/ACCLPP/blood_lead_levels.			

Maine Shared Community Health Needs Assessment Data Sources			
2015			
Indicator	Data Source	Year(s)	Other Notes
			htm(http://www.cdc.gov/nceh/lead/acclpp/blood_le ad_levels.htm
Children with unconfirmed elevated blood lead levels (% among those screened)	Maine CDC Lead Program	2009-2013	In 2012, CDC defined a reference level of 5 micrograms per deciliter (µg/dL) to identify children with elevated blood lead levels. These children are exposed to more lead than most children. For more information, visit: www.cdc.gov/nceh/lead/ACCLPP/blood_lead_levels. htm(http://www.cdc.gov/nceh/lead/acclpp/blood_le ad_levels.htm
Homes with private wells tested for arsenic	BRFSS	2009, 2012	Data are weighted to the household. At the county level, 9.7%-32.2% of those surveyed did not know whether they had tested their well water for arsenic.
Lead screening among children age 12-23 months	Maine CDC Lead Program	2009-2013	A blood lead test is considered a "screening test" only when a child has no prior history of a confirmed elevated blood lead level.
Lead screening among children age 24-35 months	Maine CDC Lead Program	2009-2013	A blood lead test is considered a "screening test" only when a child has no prior history of a confirmed elevated blood lead level.
Immunization			
Adults immunized annually for influenza	BRFSS	2011-2013	Adults who have had either a seasonal flu shot or a seasonal flu vaccine that was sprayed in your nose during the past 12 months.
Adults immunized for pneumococcal pneumonia (ages 65 and older)	BRFSS	2011-2013	Risk factor for adults aged 65 or older that have ever had a pneumonia shot.
Immunization exemptions among kindergarteners for philosophical reasons	Maine Immunization Program	2015	Available from: http://www.maine.gov/dhhs/mecdc/infectious- disease/immunization/publications/index.shtml
Two-year-olds up to date with "Series of Seven Immunizations" 4- 3-1-3-3-1-4	Maine Immunization Program	2015	The Maine Immunization Program conducts an annual immunization assessment on January 1 of each calendar year that includes all 2-year-olds in the State of Maine immunization registry, ImmPact, associated to a practice that enters client specific data. These assessments follow the standard Centers for Disease Control and Prevention childhood assessment criteria of 24-35 months of age immunized as of 24 months for the 4 DTaP (Diphtheria, Tetanus, Polio): 3 IPV (Polio): 1 MMR (Measles, Mumps, Rubella): 3 Hib (Haemophilus influenza type B): 3 HepB (Hepatitis B):1 Var (Varicella):4 PCV (Pneumococcal Conjugate) schedule.
Infectious Disease	·		
Hepatitis A (acute) incidence per 100,000 population	Maine Infectious Disease Surveillance System (MIDSS)	2014	Defined as the number of new infections during 2014.

# Maine Shared Community Health Needs Assessment Data Sources

2015				
Indicator	Data Source	Year(s)	Other Notes	
Hepatitis B (acute) incidence per 100,000 population	MIDSS	2014	Defined as the number of new infections during 2014.	
Hepatitis C (acute) incidence per 100,000 population	MIDSS	2014	Defined as the number of new infections during 2014.	
Incidence of past or present hepatitis C virus (HCV) per 100,000 population	MIDSS	2014	New diagnoses, regardless of when infection occurred or stage of disease at diagnosis.	
Incidence of newly reported chronic hepatitis B virus (HBV) per 100,000 population	MIDSS	2014	New diagnoses, regardless of when infection occurred or stage of disease at diagnosis.	
Lyme disease incidence per 100,000 population	MIDSS	2014	Defined as the number of new infections during 2014.	
Pertussis incidence per 100,000 population	MIDSS	2014	Incidence is defined as the number of new infections during 2014.	
Tuberculosis incidence per 100,000 population	MIDSS	2014	New diagnoses, regardless of when infection occurred or stage of disease at diagnosis.	
STD/HIV				
AIDS incidence per 100,000 population	Maine CDC HIV Program	2014	Incidence is defined as the number of new infections during 2014.	
Chlamydia incidence per 100,000 population	Maine CDC STD Program	2014	Incidence is defined as the number of new infections during 2014.	
Gonorrhea incidence per 100,000 population	Maine CDC STD Program	2014	Incidence is defined as the number of new infections during 2014.	
HIV incidence per 100,000 population	Maine CDC HIV Program	2014	Incidence is defined as the number of new infections during 2014.	
HIV/AIDS hospitalization rate per 100,000 population	MHDO	2011	DRG-MDC 25	
Syphilis incidence per 100,000 population	Maine CDC STD Program	2014	Incidence is defined as the number of new infections during 2014.	
Intentional Injury				
Domestic assaults reports to police per 100,000 population	Maine Dept. of Public Safety	2013	All offenses of assault between family or household members are reported as domestic assault.	
Firearm deaths per 100,000 population	Maine CDC Vital Records	2009-2013	ICD-10 W32-W34 ,X72-X74, X93-X95, Y22- Y24, Y350 or U014.	
Intentional self-injury (Youth)	MIYHS	2013	High school students who have ever done something to purposely hurt themselves without wanting to die, such as cutting or burning themselves on purpose.	
Lifetime rape/non-consensual sex (among females)	BRFSS	2012	Females who have ever had sex with someone after they said or showed that they didn't want them to or without their consent.	
Nonfatal child maltreatment per 1,000 population	Child Maltreatment Report ACYF	2013	Rates are unique child victims per 1,000 population under age 18.	
Reported rape per 100,000 population	Maine Dept. of Public Safety	2013	Includes rape by force and attempted forcible rape. Excludes carnal abuse without force (statutory rape) and other sex offenses.	
Suicide deaths per 100,000 population	Maine CDC Vital Records	2009-2013	ICD-10 U03 X60-X84 or Y87.0	

#### Maine Shared Community Health Needs Assessment Data Sources 2015 Indicator **Data Source** Year(s) **Other Notes** Violence by current or former Females who have experienced physical violence or BRFSS 2012 intimate partners in past 12 had unwanted sex with a current or former intimate months (among females) partner within the past 12 months. Reported violent crime offenses. Violent crime Violent crime rate per 100,000 Maine Dept. of 2013 includes murder, rape, robbery and aggravated population Public Safety assault. Adults reporting they always use seatbelts when they Always wear seatbelt (Adults) BRFSS 2013 drive or ride in a car. Always wear seatbelt (High School High School students who report they always wear a MIYHS 2013 Students) seatbelt when riding in a vehicle. Emergency department visits by Maine residents at Maine acute care hospitals that did not end with the patient being admitted to that hospital as an inpatient, for which the principal diagnosis is an Traumatic brain injury related injury (ICD 9 CM 800-909.2, 909.4, 909.9-994.9, emergency department visits (all MHDO 2011 995.5-995.59 or 995.80-995.85) or any external intents) per 10,000 population cause of injury code is ICD 9 CM E800-E869, E880-E929 or E950-E999, and the principal or any other diagnosis is ICD-9-CM 800.00-801.99, 803.00-804.99, 850.0-850.9, 851.00-854.19, 950.1-950.3, 959.01 or 995.55. Unintentional and undetermined Maine CDC Vital Deaths of Maine residents for which the underlying 2009-2013 intent poisoning deaths per Records cause of death is ICD-10 X40-X49 or Y10-Y19. 100,000 population Maine CDC Vital Unintentional fall related deaths Deaths of Maine residents for which the underlying 2009-2013 per 100,000 population Records cause of death is ICD-10 W00-W19. Unintentional fall related injury Unintentional fall-related injury ED Visits are defined 2011 emergency department visits per MHDO as ED Visits in which external cause of injury was 10,000 population coded as ICD--9CM E880-E886 or E888. Deaths of Maine residents for which the underlying cause of death is ICD-10 V02-V04 (.1, .9), V09.2, V12-Unintentional motor vehicle traffic Maine CDC Vital V14 (.3-.9), V19 (.4-.6), V20-V28 (.3-.9), V29 (.4-.9), 2009-2013 crash related deaths per 100,000 Records V30-V39 (.4-.9), V40-V49 (.4-.9), V50-V59 (.4-.9) , V60population V69 (.4-.9), V70-V79 (.4-.9) ,V80 (.3-.5), V81.1 ,V82.1, V83-V86 (.0-.3) ,V87 (.0-.8) or V89.2." **Occupational Health** Includes self-employed workers, owners of Deaths from work-related injuries Maine Dept. of unincorporated businesses and farms, paid and 2013 (number) Labor unpaid family workers, members of partnerships and may include owners of incorporated businesses. Includes both injuries that required days away from Nonfatal occupational injuries U.S. Bureau of work and those that required job transfer or 2013 (number) Labor Statistics restriction. Data do not reflect the relative FTEs worked by the various groups of employees.

# Maine Shared Community Health Needs Assessment Data Sources

2015				
Indicator	Data Source	Year(s)	Other Notes	
Adults who have ever had depression	BRFSS	2011-2013	Adults who have ever been told by a doctor or other healthcare provider that they have a depressive disorder.	
Adults with current symptoms of depression	BRFSS	2011-2013	Indicator of current depression coded using two items from the PHQ-2 depression screener.	
Adults currently receiving outpatient mental health treatment	BRFSS	2011-2013	Adults now taking medicine or receiving treatment from a doctor for any type of mental health condition or emotional problem.	
Co-morbidity for persons with mental illness	BRFSS	2011, 2013	Adults with current symptoms of depression from the PHQ-2 depression screener with 3 or more chronic conditions.	
Mental health emergency department rates per 100,000 population	MHDO	2011	ICD-9 CM- 209-302, 306-319, which exclude substance use related disorders.	
Sad/hopeless for two weeks in a row (High School Students)	MIYHS	2013	During the past 12 months, did you ever feel so sad or hopeless almost every day for two weeks or more in a row that you stopped doing some usual activities? Percentage of students who answered "Yes".	
Seriously considered suicide (High School Students)	MIYHS	2013	During the past 12 months, did you ever seriously consider attempting suicide? Percentage of students who answered "Yes".	
Physical Activity, Nutrition and Weig	ht			
Fewer than two hours combined screen time (High School Students)	MIYHS	2013	Percentage of students watching 2 or fewer hours of combined screen time (tv, video games, computer) per day on an average school day.	
Fruit and vegetable consumption (High School Students)	MIYHS	2013	Percentage of students who drank 100% fruit juice, ate fruit and/or ate vegetables five or more times per day during the past seven days.	
Fruit consumption among Adults 18+ (less than one serving per day)	BRFSS	2013	Adults with less than one serving per day of fruits or fruit juice.	
Met physical activity recommendations (Adults)	BRFSS	2013	Adults who reported doing enough physical activity to meet the aerobic and strengthening recommendations.	
Physical activity for at least 60 minutes per day on five of the past seven days (High School Students)	MIYHS	2013	Percentage of students who were physically active for a total of at least 60 minutes per day on five of the past seven days.	
Sedentary lifestyle – no leisure- time physical activity in past month (Adults)	BRFSS	2011-2013	Adults reporting that during the past month, other than their regular job, they did not participate in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise.	
Soda/sports drink consumption (High School Students)	MIYHS	2013	Percentage of students who drank at least one can, bottle, or glass of soda, sports drink, energy drink, or other sugar-sweetened beverage such as Gatorade, Red Bull, lemonade, sweetened tea or coffee drinks, flavored milk, Snapple, or Sunny Delight (Not counting diet soda, other diet drinks, or 100% fruit juice.) per day during the past week.	

Maine Shared Community Health Needs Assessment Data Sources				
2015				
Indicator	Data Source	Year(s)	Other Notes	
Vegetable consumption among Adults 18+ (less than one serving per day)	BRFSS	2013	Adults with less than one serving per day of vegetables.	
Obesity (Adults)	BRFSS	2013	Adults with a BMI of 30 or more.	
Obesity (High School Students)	MIYHS	2013	Percentage of students who were obese (i.e., at or above the 95th percentile for body mass index, by age and sex) SELF-REPORTED HEIGHT/WEIGHT.	
Overweight (Adults)	BRFSS	2013	Adults with a BMI between 25.0 and 29.9.	
Overweight (High School Students)	MIYHS	2013	Percentage of students who were overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex) SELF-REPORTED HEIGHT/WEIGHT.	
Pregnancy and Birth Outcomes				
Children with special health care needs	National Survey of Children with Special Health Care Needs	2011-2012	Survey respondents who reported that their child has a special health care need.	
Infant deaths per 1,000 live births	Maine CDC Vital Records	2003-2012	Number of babies who died before their first birthday per 1,000 live births. Average annual number of infant deaths and infant mortality rate might be slightly underestimated due to possible missing out-of-state deaths of Maine infants in 2010.	
Live births for which the mother received early and adequate prenatal care	Maine CDC Vital Records	2010-2012	Defined as an adequate or adequate-plus rating on the Kotelchuck Adequacy of Prenatal Care Utilization Index.	
Live births to 15-19 year olds per 1,000 population	Maine CDC Vital Records	2010-2012	Defined as the number of live births among 15- to 19-year-old Maine women per 1,000 population.	
Low birth weight (<2500 grams)	Maine CDC Vital Records	2010-2012	Low birth weight defined as less than 2500 grams.	
Substance and Alcohol Abuse				
Alcohol-induced mortality per 100,000 population	Maine CDC Vital Records	2009-2013	ICD-10 - E24.4 , F10, G31.2, G62.1, G72.1, I42.6, K29.2, K70, K85.2, K86.0, R78.0, X45, X65 or Y15	
Binge drinking of alcoholic beverages (High School Students)	MIYHS	2013	During the past 30 days, on how many days did you have 5 or more drinks of alcohol in a row, that is, within a couple of hours? Percentage of students who answered at least 1 day.	
Binge drinking of alcoholic beverages (Adults)	BRFSS	2011-2013	Risk factor for binge drinking where binge drinking is defined as having 5 or more drinks on 1 occasion for men and 4 or more drinks on 1 occasion for women.	
Chronic heavy drinking (Adults)	BRFSS	2011-2013	At risk for heavy alcohol consumption (greater than two drinks per day for men and greater than one drink per day for women).	
Drug-affected baby referrals received as a percentage of all live births	OCFS Maine Automated Child Welfare Information System	2014	This measure reflects the number of infants born in Maine where a healthcare provider reported to OCFS that there was reasonable cause to suspect the baby may be affected by illegal substance abuse or demonstrating withdrawal symptoms resulting from	

Maine Shared Community Health Needs Assessment Data Sources				
2015				
Indicator	Data Source	Year(s)	Other Notes	
			prenatal drug exposure or who have fetal alcohol spectrum disorders.	
Drug-induced mortality per 100,000 population	CDC Wonder	2009-2013	The population figures for year 2013 are bridged- race estimates of the July 1 resident population, from the Vintage 2013 postcensal series released by NCHS on June 26, 2014.	
Emergency medical service overdose response per 100,000 population	Maine Emergency Medical Services	2014	Includes overdoses from drugs/medication, alcohol and inhalants.	
Opiate poisoning (ED visits) per 100,000 population	MHDO	2009-2011	ICD-9 - 9650, 96500, 96501, 96502, 96509	
Opiate poisoning (hospitalizations) per 100,000 population	MHDO	2009-2011	ICD-9 - 9650, 96500, 96501, 96502, 96509	
Past-30-day alcohol use (High School Students)	MIYHS	2013	During the past 30 days, on how many days did you have at least one drink of alcohol? Percentage of students who answered at least 1 day.	
Past-30-day inhalant use (High School Students)	MIYHS	2013	During the past 30 days, how many times did you sniff glue, breathe the contents of aerosol spray cans, or inhale any paints or sprays to get high? Percentage of students who answered at least 1 time.	
Past-30-day marijuana use (Adults)	BRFSS	2011-2013	During the past 30 days, have you used marijuana?	
Past-30-day marijuana use (High School Students)	MIYHS	2013	During the past 30 days, how many times did you use marijuana? Percentage of students who answered at least 1 time.	
Past-30-day nonmedical use of prescription drugs (Adult)	BRFSS	2011-2013	Adults who used prescription drugs that were either not prescribed and/or not used as prescribed in order to get high at least once within the past 30 days.	
Past-30-day nonmedical use of prescription drugs (High School Students)	MIYHS	2013	During the past 30 days, how many times did you take a prescription drug (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax) without a doctor's prescription? Percentage of students who answered at least 1 time.	
Prescription Monitoring Program opioid prescriptions (days supply/pop)	Prescription Monitoring Program	2014-2015	Presented as Days Supply/Population, which is the total days of supply of medication divided by the overall population.	
Substance-abuse hospital admissions per 100,000 population	MHDO	2011	DRG-MDC 20	
Tobacco Use				
Current smoking (Adults)	BRFSS	2011-2013	Adults that reported having smoked at least 100 cigarettes in their lifetime and currently smoke.	
Current smoking (High School Students)	MIYHS	2013	During the past 30 days, on how many days did you smoke cigarettes? Percentage of students who answered at least 1 day.	
Current tobacco use (High School Students)	MIYHS	2013	Percentage of students who smoked cigarettes or cigars or used chewing tobacco, snuff, or dip on one or more of the past 30 days. (Note: Reports read	

Maine Shared Community Health Needs Assessment Data Sources			
2015			
Indicator	Data Source	Year(s)	Other Notes
			"Percentage of students who smoked cigarettes and/or cigars and/or used chewing tobacco, snuff, or dip on one or more of the past 30 days").
Secondhand smoke exposure (Youth)	MIYHS	2013	Percentage of students who were in the same room with someone who was smoking cigarettes at least 1 day during the past 7 days.

#### We wish to thank many people who provided input to this report.

Funding Partners:

Peter E. Chalke, Central Maine HealthCare, President and CEO M. Michelle Hood, FACHE, EMHS President and CEO Chuck Hays, MaineGeneral Health, CEO and President William L. Caron, Jr., MaineHealth, President Mary C. Mayhew, Maine DHHS, Commissioner

Market Decisions/Hart Consulting, Inc. Research Team: Patrick Madden, MBA Patricia Hart, MS, GC-PH John Charles Jennifer MacBride Bethany Porter Kelly MacGuirl, MSc

University of Southern Maine, Muskie School of Public Service, Epidemiologist Team:

Crystal Cushman Zachariah Croll Kathy Decker Pamela Foster Albert Alison Green-Parsons Sara Huston Jennifer Lenardson Erika Lichter Cindy Mervis Alexandra Nesbitt Donald Szlosek Finn Teach Denise Yob Erika Ziller

Maine SHNAPP Steering Committee:

Nancy Birkhimer - Director, Performance Improvement, Maine CDC, Maine DHHS Deborah Deatrick - Senior Vice President, Community Health Improvement, MaineHealth Doug Michael - Chief Community Health & Grants Officer, Eastern Maine Healthcare Systems Natalie Morse - Director of the Center for Prevention and Healthy Living, MaineGeneral Cindie Rice - Director of Community Health, Wellness and Cardiopulmonary Rehab, Central Maine Medical Center Maine SHNAPP Metrics Subcommittee: Nancy Birkhimer, Maine CDC, Maine DHHS Sean Cheetham. Central Maine Medical Center Tim Cowan, MaineHealth Ron Deprez, University of New England Brent Dubois, Eastern Maine Healthcare Systems Charles Dwyer, Maine Health Access Foundation Jayne Harper, SHNAPP Staff (MaineGeneral Health) Rebecca Kingsbury, MaineGeneral Health Jean Mellett, Eastern Maine Healthcare Systems Natalie Morse, MaineGeneral Health Jeb Murphy, Maine Primary Care Association Lisa Nolan, Maine Health Management Coalition Rebecca Parent, Eastern Maine Healthcare Systems Sandra Parker, Maine Hospital Association Cindie Rice, Central Maine Medical Center Toho Soma, Portland Public Health Division Jenn Yurges, MaineGeneral Health Maine SHNAPP Community Engagement Subcommittee: Nancy Birkhimer, Maine CDC, Maine DHHS

Andy Coburn, University of Southern Maine, Muskie School Charles Dwyer, Maine Health Access Foundation Deb Erickson-Irons, York Hospital Joanne Fortin, Northern Maine Medical Center Nicole Hammar, Eastern Maine Healthcare Systems Jayne Harper, SHNAPP Staff (MaineGeneral Health) Elizabeth Keene, St. Mary's Regional Medical Center Celine Kuhn, MaineHealth Joy Leach, MaineGeneral Health Christine Lyman, Maine CDC, Maine DHHS Becca Matusovich, formerly Maine CDC, Maine DHHS Doug Michael, Eastern Maine Healthcare Systems Natalie Morse, MaineGeneral Health Jeb Murphy, Maine Primary Care Association Cindie Rice, Central Maine Medical Center Toho Soma, Portland Public Health Division Paula Thomson, Maine CDC, Maine DHHS

### Collaborating Organizations for SHNAPP Implementation:

Bangor Public Health and Community Services Maine Health Access Foundation Maine Health Management Coalition Maine Hospital Association Maine Office of Substance Abuse and Mental Health Services Maine Primary Care Association Portland Public Health Division St. Mary's Regional Medical Center Statewide Coordinating Council for Public Health University of New England University of Southern Maine, Maine Public Health Institute at the Muskie School

Maine Department of Health and Human Services Review Team: Ken Albert, Maine CDC Director and Chief Operating Officer Sheryl Peavey, DHHS, Strategic Reform Coordinator Jay Yoe, Director, DHHS Office of Continuous Quality Improvement

District Public Health Liaisons:

Aroostook: Stacy Boucher Central: Paula Thomson Cumberland: Becca Matusovich, formerly Maine CDC, Maine DHHS Cumberland: Adam Hartwig, acting Downeast: Alfred May MidCoast: Carrie McFadden Penquis: Jessica Fogg Wabanaki: Kristi Ricker and Sandra Yarmal Western: Jamie Paul York: Adam Hartwig