

Testing Schedule/Result Availability for Tests Not Performed Daily (Subject to change during holiday weeks)

| COAGULATION | |
|---|---|
| TEST | TESTING SCHEDULE/RESULT AVAILABILITY |
| DRVVT screen and confirm/ Silica Clot Time | T, Th |
| AT3, Protein C and S | W |
| PF4 for HIT | Sun - F |
| Platelet Aggregation/vonWillebrand Panel | W |
| Ristocetin Cofactor (vonWillebrand Factor Activity) | W (Stat by request) |
| Sickle Cell | M, W, F |
| GENERAL LABORATORY | |
| ANA | M-F |
| Immunofixation Electrophoresis (serum, urine) | M-F (reported within 72 hours) |
| Myoglobin | M- F (available STAT) |
| Protein Electrophoresis Testing (serum, urine) | M-F (reported within 72 hours) |
| Referral Test | 2 days or more (sent to off-site laboratory) |
| MICROBIOLOGY | |
| AFB | Cultures held 6 weeks |
| Anticardiolipin IgG and IgM | M |
| BP PCR (B. pertussis) | M-F |
| Bacterial Culture and Sensitivity | Clinically significant information is often available within 24 |
| | hours. Final results vary with culture types. |
| Celiac TTG Panel | M, W, F |
| EBV Antibodies | T, F |
| FAC2PCR (Factor II) | 2 days or more (sent to off-site laboratory) |
| FAC5PCR (Factor V) | 2 days or more (sent to off-site laboratory) |
| HCV RNA Quantitative | W |
| HIV Antibody | M, W, F |
| HIV Q PCR | W |
| H.pylori stool antigen | M, W, F |
| HSV PCR (non-CSF) | M, W, F |
| HSV SER (Type 1 & 2 Serology) | T, F |
| Lyme IgG/IgM Antibody Screen | M, W, F (daily during peak season) |
| (reflexed to WB if positive) | |
| Mumps IgG Immune Status | M, W, F |
| Rubella IgG Immune Status | M,W, F |
| Rubeola IgG Immune Status | M,W, F |
| TTG-A and TTG-G | M, W, F |
| Varicella Zoster IgG Immune Status | M, W, F |
| VZV PCR (non-CSF) | 2 days or more (sent to off-site laboratory) |
| TOXICOLOGY | |
| Cyclosporine | M – F |
| Tacrolimus | M – F |